

**ADVENTIVE APHIDS (HEMIPTERA: APHIDIDAE)
OF AMERICA NORTH OF MEXICO**

ROBERT G. FOOTIT, SUSAN E. HALBERT, GARY L. MILLER, ERIC MAW,
AND LOUISE M. RUSSELL¹

(RGF, EM) Agriculture and Agri-Food Canada, Canadian National Collection of Insects, Ottawa, ON, K1A 0C6 Canada (e-mail: footitrg@agr.gc.ca; mawe@agr.gc.ca); (SEH) Florida Department of Agriculture and Consumer Services, Division of Plant Industry, Gainesville, FL 32614, U.S.A. (e-mail: halbers@doacs.state.fl.us); (GLM, LMR) Systematic Entomology Laboratory, Plant Science Institute, Agricultural Research Service, U.S. Department of Agriculture, Beltsville, MD 20705, U.S.A. (e-mail: gmiller@sel.barc.usda.gov)

Abstract. — We provide a compilation of 262 species of aphids that are considered as adventive to North America north of Mexico. Included for each species, where applicable, is reference to: the location and date of introduction of the first North American record; pest status in North America; principal economic hosts; and biogeographical origin. Information is also provided for species whose presence in North America is considered erroneous or questionable and for those species that are considered Holarctic or Beringian.

Key Words: Aphidoidea, plant lice, plant pathogens, invasive species, biological control, quarantine, agriculture, forestry, horticulture, ornamentals, fruit trees

In an era of a global economy and commerce, the increase of adventive species represents both domestic and international concerns. If non-indigenous species become pests, the economic consequences include loss of production, diminished quality, production cost increases, flexibility decreases in production/management decisions, increased risk of human disease, and damaging environmental and aesthetic effects (Huber et al. 2002). Various insects of agricultural concern have been identified as potentially damaging if introduced or

reintroduced into the United States (Huber et al. 2002).

Many aphid species have been added to the North American fauna since colonization by Europeans, and some of these have become serious pests. In addition to direct feeding damage, aphids constitute one of the largest groups of known plant-virus vectors (Chan et al. 1991). Aphids are primarily controlled through pesticide applications, adding to crop production costs, and having environmental implications associated with increased pesticide use. The effects of some non-indigenous aphids in North America have been staggering. For example, by the early 1990's, the Russian wheat aphid, *Diur-*

¹ It is noteworthy that Louise Russell, a pioneer woman entomologist and specialist in the Sternorrhyncha, continues to publish at the age of 101.

aphis noxia (Kurdjumov), was responsible for \$500–900 million cumulative losses in the western U.S. (Bernal et al. 1993, Morrison and Peairs 1998). The appearance of the brown citrus aphid, *Toxoptera citricida* (Kirkaldy), in Florida in 1995 had a major impact on the U.S. citrus industry (Halbert and Brown 1996). More recently, losses related to the soybean aphid, *Aphis glycines* Matsumura, have been estimated at about \$120 million in 2003 for Minnesota alone (Anonymous 2005).

The origins of this work actually began in part in the 1970's when one of us (LMR) contributed data on invasive species in the United States (e.g., Sailer 1978). Nearly 30 years later, adventive insects are still a major concern for agriculture in North America. The objectives of this paper are to: 1) list the adventive aphid species (excluding adelgids and phylloxerans) of North America and record data on first detection date (along with validation source), principal hosts, and zoogeographic area of origin; 2) analyze these data and determine if patterns exist; and 3) compare these data to other sternorrhynchan groups.

MATERIALS AND METHODS

We use the term adventive species as defined by D.R. Miller et al. (2002), i.e., for those species that are non-native to North America regardless of economic harm. Other terms such as invasive, alien, exotic, and immigrant have sometimes been used interchangeably (see Wheeler and Hoebeke (2001) and Miller et al. (2005) for discussion). Using Smith and Parron's (1978) list of North American aphids as the base, we compiled a new list for North American north of Mexico. We exclude species that have been intercepted at ports-of-entry or discovered in nurseries or greenhouses and subsequently eradicated. With the exception of *Aphis chloris* Koch, a bio-control agent for *Hypericum perforatum*

(Harris 2003), no aphid species has been purposely introduced.

Geographic origins are given in general terms only, relying heavily on Blackman and Eastop (1994, 2000), supplemented by other references, as noted in Table 1, and by opinion of the authors. The origins of a few cosmopolitan species are unknown. It is not always readily apparent whether a species is adventive or native; some species we have listed as adventive may in fact be naturally Holarctic in distribution and vice versa.

For determining the earliest record of the North American adventive aphids, we relied on published records and on specimens examined in The Canadian National Collection of Insects, Nematodes and Arachnids (CNC) and the Edith Patch Collection (CNC-Patch) housed at CNC, Ottawa, ON, Canada; the MacGillivray Collection (ACNB), Agriculture and Agri-Food Canada, Fredericton, NB, Canada; The Florida State Collection of Arthropods (FSCA), Gainesville, FL, U.S.A.; and The National Entomological Collection of the National Museum of Natural History (USNM), Beltsville, MD, U.S.A. In addition, Asa Fitch's aphid species notes that are currently housed with the USNM aphid collection at Beltsville, MD, were also examined. This collection of written records and observations from the 1840's–1870's provide some of our earliest glimpses into the North American aphid fauna. While there are some questions concerning Fitch's "species concept" versus current thinking for some of his included species, there is little confusion given the morphology and associated host plants for others. His reference to the occurrence in 1791 of the cabbage aphid, *Brevicoryne brassicae* (L.) is possibly the earliest record of an adventive aphid for North America.

Nomenclature followed herein reflects that of Remaudière and Remaudière

Table 1. Introduced aphid species in Canada and continental United States of America, giving earliest record with source of record, geographic origin, pest status, and economic host species. Aphid names follow Remaudière and Remaudière (1997). Earliest records are based on literature references and on specimens in the following collections: USNM — National Museum of Natural History; CNC — Canadian National Collection of Insects, Arachnids and Nematodes; FSCA — Florida State Collection of Arthropods; ACNB — MacGillivray collection, Agriculture and Agri-Food Canada, Fredericton, NB; CNC-Patch — Edith Patch collection, housed at CNC; EMEC — Essig Museum of Entomology Collection, University of California, Berkeley.

Introduced Species	Earliest N.A. record (1) (Collection or Reference)	Geographic Origin/ Reference (2)	Status in North America/ Reference (2)	North American Economic Host(s)
<i>Acyrtosiphon caraganae</i> (Cholodkovsky)	SK, 1940 (CNC)	Palaearctic	pest	<i>Caragana arborescens</i>
<i>Acyrtosiphon kondoi</i> Shinji	CA, 1974 (Kono 1977)	Palaearctic (e Asia)	pest Halbert et al. 2000	<i>Medicago sativa</i>
<i>Acyrtosiphon lactucae</i> (Passerini)	MD, 1943 (USNM)	Palaearctic (Europe)		<i>Lactuca sativa</i>
<i>Acyrtosiphon malvae</i> (Mosley) (3)	NE, 1891 (USNM)	Palaearctic (Europe)		<i>Geranium, Pelargonium</i>
<i>Acyrtosiphon pisum</i> (Harris)	KS, 1877 (Harper et al. 1978)	Palaearctic (Europe)	pest	Leguminosae, incl. <i>Pisum</i> , <i>Medicago</i>
<i>Aloephagus myersi</i> Essig	CA, 1939 (Essig 1950)	Afrotropical	minor pest Essig 1950, regulatory issue	<i>Aloe</i> (greenhouse)
<i>Amphorophora ampullata</i> Bucktön	MN, 1887 (Oestland 1887)	Palaearctic (Europe)		
<i>Anoecia corni</i> (Fabricius)	IA, 1897 (USNM)	Palaearctic (Europe)		roots of Gramineae, incl. cereals
<i>Aphidounguis mali</i> Takahashi (4)	NC, 1999 (CNC)	Palaearctic (e Asia)		
<i>Aphis amaranthi</i> Holman	FL, 1990 (Halbert et al. 2000)	Neotropical (Cuba)	potential pest Halbert et al. 2000	edible <i>Amaranthus</i>
<i>Aphis callunae</i> Theobald	BC, 1974 (Forbes and Chan 1976)	Palaearctic (Europe)		
<i>Aphis chloris</i> Koch	BC, 1980 (Harris 2003)	Palaearctic (Europe)	Released as weed biological control agent	<i>Hypericum perforatum</i>
<i>Aphis craccae</i> Linnaeus	BC, NB, ME, 1965 (CNC, ACNB)	Palaearctic		<i>Vicia cracca</i>
<i>Aphis craccivora</i> Koch (5)	NE, 1890 (USNM)	Palaearctic		polyphagous, with preference for legumes
<i>Aphis cytisorum</i> Hartig	DC, 1906 (USNM)	Palaearctic (Europe)		<i>Cytisus scoparius</i>
<i>Aphis epilobii</i> Kaltenbach	CO, 1915 (USNM)	Palaearctic (Europe)		
<i>Aphis euphorbiae</i> Kaltenbach (6)	CA, 1933 (USNM)	Palaearctic (Europe)		
<i>Aphis fabae</i> Scopoli (7)	NY, 1869 (Fitch 1870 [as <i>A. rumicis</i> L.])	Palaearctic (Europe)	pest	polyphagous
<i>Aphis genistae</i> Scopoli	MA, 1921 (USNM)	Palaearctic (Europe)		
<i>Aphis glycines</i> Matsumura	IA, IL, IN, KY, MI, MN, MO, OH, WI, WV, 2000 (Voegtlin 2000)	Oriental and Palaearctic (East Asia)	major pest Ragsdale et al. 2004	<i>Glycine max</i>
<i>Aphis gossypii</i> Glover	NY, 1852 (Russell 1968)	Palaearctic	major pest	polyphagous

Table 1. Continued.

Introduced Species	Earliest N.A. record (1) (Collection or Reference)	Geographic Origin/ Reference (2)	Status in North America/ Reference (2)	North American Economic Host(s)
<i>Aphis hederæ</i> Kaltenbach	CA, 1910 (Essig 1910)	Palearctic (Europe)	minor pest	<i>Hedera helix</i>
<i>Aphis idæi</i> van der Goot	BC, 1951 (Forbes et al. 1973)	Palearctic (Europe)		
<i>Aphis nasturtii</i> Kaltenbach	NE, 1894 (USNM)	Palearctic	pest	<i>Solanum tuberosum</i>
<i>Aphis nepetæ</i> Kaltenbach	ID, 1957 (Gittins et al. 1976)	Palearctic		
<i>Aphis nerii</i> Boyer de Fonscolombe	IL, 1879 (Thomas 1879)	Palearctic (e Asia)	pest	ornamentals (flowers)
<i>Aphis polygonata</i> (Nevsky) (8)	IL, 1977 (FSCA)	Palearctic (Central Asia)		
<i>Aphis pomi</i> De Geer	PA, 1844 (Haldeman 1844)	Palearctic (Europe)	pest	<i>Malus pumila</i> , <i>Pyrus communis</i> , <i>Cydonia oblonga</i>
<i>Aphis praeterita</i> Walker	ME, 1918 (Patch 1918)	Palearctic (Europe)		
<i>Aphis rumicis</i> Linnaeus (7)	Unknown (7)	Palearctic (Europe)		
<i>Aphis salicariæ</i> Koch	ON, 1952 (CNC)	Palearctic (Europe)		
<i>Aphis sambuci</i> Linnaeus	NY, 1847 (Fitch notes [as <i>A. sambucifoliæ</i> Fitch])	Palearctic (Europe)		
<i>Aphis sedi</i> Kaltenbach	ME, 1911 (Patch 1911)	Palearctic (Europe)		
<i>Aphis spiræcola</i> Patch	CO, 1907 (Gillette 1910)	Palearctic or Oriental	major pest	polyphagous
<i>Aphis taraxicola</i> (Börner) (9)	QC, 1983 (Cloutier et al. 1986)	Palearctic		
<i>Aphis thalictri</i> Koch	BC, 1999 (CNC)	Palearctic (Europe)		
<i>Aphis umbrella</i> (Börner)	NE, SD, 1891 (USNM [as <i>A. maliae</i> Koch])	Palearctic (Europe)		
<i>Aphis urticata</i> Gmelin	PA, 1913 (Olsen 1921)	Palearctic (Europe)		
<i>Aphis vaccini</i> (Börner)	BC, 2000 (CNC)	Palearctic (Europe)		
<i>Aploneura lentisci</i> (Passerini)	IL, 1879 (Thomas 1879 [as <i>Rhizobius poæ</i> Thomas])	Palearctic (Mediterranean)		Gramineae, including <i>Triticum</i>
<i>Asiphonella dactylonii</i> Theobald	AZ, 1947 (USNM)	Palearctic (e Asia)	potential pest	<i>Cynodon dactylon</i>
<i>Aspidaphis adjuvans</i> (Walker)	CO, 1907 (Gillette 1917b)	Palearctic (Europe)		
<i>Atarsaphis agrifoliæ</i> (Ferris)	CA, 1921 (Ferris 1921)	Palearctic (e Asia)		
<i>Atheroides serrulatus</i> Haliday	QC, 1966 (CNC)	Palearctic	minor pest	Gramineae, incl. cereals
<i>Aulacorthum circumflexum</i> (Buckton)	CO, 1907 (Gillette 1908)	Palearctic (e Asia) or Oriental	pest	polyphagous
<i>Aulacorthum solani</i> (Kaltenbach)	NY, 1879 (Riley and Monell 1879)	Palearctic	major pest	polyphagous
<i>Betulaphis brevopilosa</i> Börner	BC, 1959 (Forbes et al. 1974)	Palearctic (Europe)		

Table 1. Continued.

Introduced Species	Earliest N.A. record (1) (Collection or Reference)	Geographic Origin/ Reference (2)	Status in North America/ Reference (2)	North American Economic Host(s)
<i>Betulaphis helvetica</i> Hille Ris Lambers	BC, 1926 (Glendenning 1926)	Palaearctic (Europe)		
<i>Betulaphis quadrituberculata</i> (Kaltenbach)	BC, 1946 (ACNB)	Palaearctic (Europe)		
<i>Brachycaudus helichrysi</i> (Kaltenbach)	OR, 1893 (USNM)	Palaearctic (Europe)	major pest	<i>Prunus</i> , Compositae
<i>Brachycaudus cardui</i> (Linnaeus)	NY, 1887 (USNM)	Palaearctic (Europe)	pest	<i>Prunus</i> , Compositae
<i>Brachycaudus persicae</i> (Passerini)	MD, DE, NJ, VA, MI, 1887 (Smith 1890)	Palaearctic (Europe)	pest	<i>Prunus</i>
<i>Brachycaudus rumexicolens</i> (Patch) (10)	CT 1913 (Patch 1917)	Palaearctic (Europe)		
<i>Brachycaudus schwartzi</i> (Börner)	CA, 1975 (Eastop 1975)	Palaearctic (Europe)	pest	<i>Prunus persica</i>
<i>Brachycaudus spiraeae</i> (Börner)	ON, 1956 (CNC)	Palaearctic		
<i>Brachycaudus tragopogonis</i> (Kaltenbach)	WA, 1993 (FSCA)	Palaearctic		
<i>Brachycorynella asparagi</i> (Mordvilko)	NY, NJ, 1969 (Leonard 1970, Race 1970)	Palaearctic (Europe)	locally serious pest Stoetzel 1990	<i>Asparagus officinalis</i>
<i>Brevicoryne brassicae</i> (Linnaeus)	NY, 1791 (Fitch notes)	Palaearctic	pest	<i>Brassica</i> crops
<i>Calaphis betulicola</i> (Kaltenbach)	NY, 1854 (Fitch notes)	Palaearctic (Europe)		
<i>Callipterinella calliptera</i> (Hartig)	OR, 1909 (USNM)	Palaearctic (Europe)	minor pest Hajek and Dahlsten 1986	<i>Betula</i>
<i>Callipterinella minutissima</i> (Stroyan)	BC, 1975 (Forbes and Chan 1980)	Palaearctic (Europe)	minor pest Hajek and Dahlsten 1987	<i>Betula</i>
<i>Capitophorus carduinus</i> (Walker)	GA, 1948 (USNM)	Palaearctic (Europe)	potential weed biocontrol agent Gassmann and Kok 2002	<i>Cirsium</i>
<i>Capitophorus elaeagni</i> (Del Guercio)	SD, 1892 (USNM)	Palaearctic (Europe)	Potential virus vector	
<i>Capitophorus hippophaes</i> (Walker)	DC, 1895 (USNM)	Palaearctic (Europe)		
<i>Capitophorus pakansus</i> Hottes and Frison	IL, 1929 (Hottes and Frison 1931)	Palaearctic (Europe)		
<i>Capitophorus horni</i> Börner	ME, 1951 (Corpuz- Raros and Cook 1974)	Palaearctic (Europe)		
<i>Cavariella aegopodii</i> (Scopoli)	MN, 1903 (USNM)	Palaearctic (Europe)	pest	<i>Daucus carota</i>
<i>Cavariella archangelicae</i> (Scopoli)	WA, 1953 (USNM)	Palaearctic (Europe, Iceland)		
<i>Cavariella pastinacae</i> (Linnaeus)	NY, 1855 (Fitch notes)	Palaearctic (Europe)		
<i>Cerataphis brasiliensis</i> (Hempel)	OH, 1917 (USNM)	Oriental	pest	palms
<i>Cerataphis lataniae</i> (Boisduval)	NE, 1912 (USNM)	Oriental	pest	palms
<i>Cerataphis orchidearum</i> (Westwood)	DC, 1923 (USNM)	Oriental	pest	greenhouse orchids

Table 1. Continued.

Introduced Species	Earliest N.A. record (1) (Collection or Reference)	Geographic Origin/ Reference (2)	Status in North America/ Reference (2)	North American Economic Host(s)
<i>Ceruraphis eriophori</i> (Walker)	UT, 1935 (Knowlton and Smith 1936)	Palearctic (Europe)		
<i>Chaetosiphon potentillae</i> (Walker)	NY, 1851 (Fitch notes) CO, 1916 (USNM)	Palearctic (Europe)		
<i>Chaetosiphon tetrarhodum</i> (Walker)	MN, 1907 (USNM)	Palearctic (Europe)		
<i>Chaitophorus leucomelas</i> Koch	CA, QC, 1966 (USNM, Quednau 1966)	Palearctic (Europe)		
<i>Chaitophorus populialbae</i> (Boyer de Fonscolombe)	ON, 1955 (CNC)	Palearctic (Europe)		
<i>Chromaphis juglandicola</i> (Kaltenbach)	CA, 1896 (USNM)	Palearctic (Europe)	pest Olsen 1974	<i>Juglans regia</i>
<i>Cinara cedri</i> Mimeur	NY 2004 (CNC, FSCA)	Palearctic (Europe)		<i>Cupressus</i> , <i>Juniperus</i>
<i>Cinara confinis</i> (Koch)	CO, 1920 (USNM)	Palearctic		<i>Abies</i>
<i>Cinara costata</i> (Zetterstedt)	ON, 1905 (USNM)	Palearctic		<i>Picea</i>
<i>Cinara cupressi</i> (Buckton)	CA, 1916 (Bartholomew 1932)	Palearctic		<i>Cupressus</i> , <i>Juniperus</i>
<i>Cinara fresai</i> Blanchard	FL, 1942 (Tissot 1944 [as <i>C.</i> <i>wacasassae</i>])	unknown		<i>Cupressus</i> , <i>Juniperus</i>
<i>Cinara juniperi</i> (De Geer)	NY, 1902 (USNM)	Palearctic (Europe)		<i>Juniperus</i>
<i>Cinara laricis</i> (Hartig)	MA, 1921 (USNM)	Palearctic (Europe)		<i>Larix</i>
<i>Cinara pilicornis</i> (Hartig)	ME, 1906 (Patch 1912)	Palearctic		<i>Picea</i>
<i>Cinara pinea</i> (Mordvilko)	DC, 1919 (Wilson 1919)	Palearctic		<i>Pinus</i>
<i>Cinara pruinosa</i> (Hartig)	CO, 1910 (Gillette 1917a [as <i>Lachnus</i> <i>palmerae</i>])	Palearctic		<i>Picea</i>
<i>Cinara tujafilina</i> (Del Guercio)	CA, 1914 (Davidson 1914)	unknown		
<i>Coloradoa abrotani</i> (Koch)	NB, 1964 (ACNB)	Palearctic (Central Asia)		
<i>Coloradoa absinthii</i> (Lichtenstein)	MB, BC, 1965 (CNC)	Palearctic (Central Asia)		
<i>Coloradoa achilleae</i> Hille Ris Lambers	ON, 1962 (CNC)	Palearctic (Central Asia)		<i>Achillea</i>
<i>Coloradoa artemisiae</i> (Del Guercio)	BC, 1965 (CNC)	Palearctic (Central Asia)		
<i>Coloradoa rufomaculata</i> (Wilson)	NE, 1890 (USNM)	Palearctic (Central Asia)	minor pest	chrysanthemums
<i>Coloradoa tanacetina</i> (Walker)	RI, 1974 (Smith and Parron 1978)	Palearctic (Central Asia)		
<i>Corylobium avellanae</i> (Schrank)	BC, 1983 (Forbes and Chan 1984)	Palearctic (Europe)	pest	<i>Corylus</i> <i>avellana</i>
<i>Cryptaphis poae</i> (Hardy)	MB, 1965 (Robinson and Bradley 1965)	Palearctic (Europe)		
<i>Cryptomyzus galeopsidis</i> (Kaltenbach)	ME, 1914 (Patch 1914)	Palearctic (Europe)		

Table 1. Continued.

Introduced Species	Earliest N.A. record (1) (Collection or Reference)	Geographic Origin/ Reference (2)	Status in North America/ Reference (2)	North American Economic Host(s)
<i>Cryptomyzus ribis</i> (Linnaeus)	NY, 1854 (Fitch notes)	Palaearctic (Europe)	pest	<i>Ribes</i>
<i>Ctenocallis setosus</i> (Kaltenbach)	OR, 1951 (Essig 1952)	Palaearctic (Europe)		
<i>Ctenocallis zelkowae</i> (Shinji)	CA, 1983 (Kono 1983)	Palaearctic (E Asia)	Pest Kono, 1983	<i>Zelkova</i>
<i>Decorosiphon</i> <i>corynothrix</i> Börner	NC, 1963 (Smith and Knowlton 1975)	Palaearctic (Europe)		
<i>Diuraphis frequens</i> Walker	ON, 1951 (CNC)	Palaearctic (Europe)	minor pest Halbert et al. 1992	<i>Triticum</i>
<i>Diuraphis noxia</i> (Kurdjumov)	TX, 1986 (Stoetzel 1987)	Palaearctic (Central Asia)	major pest Quisenberry and Peairs 1998	Gramineae, incl. cereals
<i>Drepanosiphum</i> <i>oregonensis</i> Granovsky	OR, 1914 (USNM)	Palaearctic		<i>Picea</i>
<i>Drepanosiphum</i> <i>platanoides</i> (Schrank)	CA, 1908 (Wilson 1909)	Palaearctic (Europe)		<i>Acer</i>
<i>Dysaphis apiifolia</i> (Theobald)	CA, 1934 (USNM)	Palaearctic	pest	Umbelliferae
<i>Dysaphis crataegi</i> (Kaltenbach) (11)	DC, 1915 (USNM)	Palaearctic	pest	Umbelliferae
<i>Dysaphis devector</i> (Walker)	QC, 1966 (Quednau 1966)	Palaearctic (Europe)		<i>Malus pumila</i>
<i>Dysaphis foeniculus</i> (Theobald) (12)	OR, 1949 (USNM)	Palaearctic (Europe)	occasional pest	Umbelliferae (including <i>Daucus</i> <i>carota</i>)
<i>Dysaphis plantaginea</i> (Passerini)	NE, 1890 (Williams 1911)	Palaearctic (Europe)	pest	<i>Malus pumila</i>
<i>Dysaphis pyri</i> (Boyer de Fonscolombe)	CO, 1978 (Smith and Parron 1978)	Palaearctic		<i>Pyrus</i> <i>communis</i>
<i>Dysaphis radicola</i> (Mordvilko)	DC, 1894 (USNM)	Palaearctic (Europe)		
<i>Dysaphis tulipae</i> (Boyer de Fonscolombe)	NH, 1903 (USNM)	Palaearctic (Europe)	pest, regulatory concern	bulbs
<i>Elatobium abietinum</i> (Walker)	CA, DC, 1915 (USNM)	Palaearctic (Europe)	pest	<i>Abies</i>
<i>Eriosoma grossulariae</i> (Schüle) (13)	ON, 1913 (CNC)	Palaearctic (Europe)		<i>Ribes</i> , <i>Ulmus</i>
<i>Eriosoma pyricola</i> Baker & Davidson	OH, 1897 (Baker and Davidson 1916)	Palaearctic (Europe)		<i>Pyrus</i> <i>communis</i>
<i>Eriosoma ulmi</i> (Linnaeus) (13)	BC, 2003 (CNC)	Palaearctic		<i>Ulmus</i> ; <i>Ribes</i>
<i>Eucallipterus tiliac</i> Schouteden	DC, 1890 (USNM)	Palaearctic (Europe)	pest	<i>Tilia</i>
<i>Eucarazzia elegans</i> (Ferrari)	CA, 1984 (Stoetzel 1985)	Palaearctic (Europe)		Labiatae
<i>Euceraphis betulae</i> (Koch) (14)	CO, 1909 (USNM)	Palaearctic (Europe)		<i>Betula</i>
<i>Euceraphis punctipennis</i> (Zetterstedt) (14)	NY, 1847 (Fitch notes)	Palaearctic (Europe)		<i>Betula</i>
<i>Eulachnus agilis</i> (Kaltenbach)	NY, 1909 (Gillette 1909)	Palaearctic (Europe)		<i>Pinus</i>
<i>Eulachnus brevipilosus</i> Börner	WA, 1954 (Hottes and Essig 1955)	Palaearctic (Europe)		<i>Pinus</i>
<i>Eulachnus rileyi</i> (Williams)	IN, 1886 (USNM)	Palaearctic (Europe)	minor pest	<i>Pinus</i>

Table 1. Continued.

Introduced Species	Earliest N.A. record (1) (Collection or Reference)	Geographic Origin/ Reference (2)	Status in North America/ Reference (2)	North American Economic Host(s)
<i>Euthoracaphis umbellariae</i> Essig	CA, 1929 (Essig 1932)	Palearctic (e Asia)		
<i>Forda formicaria</i> von Heyden	CO, 1909 (USNM)	Palearctic (Europe)		roots of Gramineae
<i>Forda marginata</i> Koch	DC, 1885 (USNM)	Palearctic (Europe)		roots of Gramineae
<i>Geoica setulosa</i> (Passerini)	ID, 1985 (FSCA)	Palearctic (Europe)	minor pest G.L. Miller et al. 2002	roots of Gramineae
<i>Geoica utricularia</i> (Passerini) (15)	IL, 1887 (USNM)	Palearctic (Europe)		roots of Gramineae, incl. cereals
<i>Geopemphigus floccosus</i> (Moreira)	FL, 1945 (USNM)	Neotropical	minor pest	roots of <i>Ipomoea batatas</i>
<i>Glyphina betulae</i> (Linnaeus)	PA, 1945 (USNM)	Palearctic (Europe)		
<i>Greenidea ficicola</i> Takahashi	FL, 2002 (Halbert 2004)	Oriental	potential pest	<i>Ficus</i>
<i>Greenidea psidii</i> van der Goot	CA, 1998 (Gill 1998a)	Oriental	minor pest, regulatory pest	<i>Myrtus</i>
<i>Hayhurstia atriplicis</i> (Linnaeus)	CA, 1887 (USNM)	Palearctic (Europe)		<i>Chenopodium</i> including <i>Ch. quinoa</i>
<i>Hyadaphis coriandri</i> (Das)	FL, 1997 (Halbert et al. 2000)	Palearctic (Central Asia)	major pest	Umbelliferae, including <i>Coriandrum</i> , <i>Foeniculum</i>
<i>Hyadaphis foeniculi</i> (Passerini)	NY, 1879 (USNM)	Palearctic (Central Asia)	minor pest	<i>Lonicera</i> ; Umbelliferae
<i>Hyadaphis tartaricae</i> (Ajzenberg)	IL, 1979 (Voegtlin 1981)	Palearctic (Central Asia)	major pest Voegtlin 1981	<i>Lonicera tatarica</i>
<i>Hyalopteroides humilis</i> (Walker)	DC, MA, 1909 (Hayhurst 1909)	Palearctic		
<i>Hyalopterus amygdali</i> (Blanchard) (16)	IL, NJ (date ?) (16) (Smith and Parron 1978)	Palearctic (e Asia)	pest	<i>Prunus</i> (s.g. <i>Amygdalus</i>)
<i>Hyalopterus pruni</i> (Geoffroy) (16)	IL, 1879 (Thomas 1879)	Palearctic (Europe)	pest	<i>Prunus</i> (s.g. <i>Prunus</i>)
<i>Hyperomyzus carduellinus</i> (Theobald)	FL, 1997 (Halbert et al. 2000)	Palearctic (e Asia)		
<i>Hyperomyzus lactucae</i> (Linnaeus)	IL, 1878 (Thomas 1879)	Palearctic (Europe)	minor pest	<i>Ribes</i>
<i>Hyperomyzus pallidus</i> Hille Ris Lambers	ME, 1914 (Hille Ris Lambers 1949)	Palearctic (Europe)		<i>Ribes</i>
<i>Hyperomyzus picridis</i> (Börner)	PA, 1942 (USNM)	Palearctic (Europe)	potential pest	<i>Ribes</i>
<i>Hyperomyzus rhinanthi</i> (Schouteden) (17)	ME, 1951 (CNC)	Palearctic (Europe)		<i>Ribes</i>
<i>Idiopterus nephrolepidis</i> Davis	IL, 1907 (Davis 1909)	Neotropical Stoetzel 1990	pest Stoetzel 1990	greenhouse ferns
<i>Jacksonia papillata</i> Theobald	ID, 1959 (USNM)	Palearctic (Europe)		
<i>Liosomaphis berberidis</i> (Kaltenbach)	NY, 1846 (Fitch notes)	Palearctic (Europe)		
<i>Lipamyzodes matthiolarum</i> (Doncaster)	WA, 1976 (Fox 1977)	Palearctic (Europe)		

Table 1. Continued.

Introduced Species	Earliest N.A. record (1) (Collection or Reference)	Geographic Origin/ Reference (2)	Status in North America/ Reference (2)	North American Economic Host(s)
<i>Lipaphis pseudobrassicæ</i> (Davis) (18)	DC, 1899 (USNM)	Paleartic (Europe)	major pest Paddock 1915	Cruciferae
<i>Longicaudus trirhodus</i> (Walker)	MA, 1908 (USNM)	Paleartic (Europe)		<i>Rosa</i>
<i>Macrosiphoniella</i> <i>abrotani</i> (Walker)	IL, 1929 (USNM)	Paleartic		
<i>Macrosiphoniella</i> <i>absinthii</i> (Linnaeus)	IL, 1879 (Thomas 1879)	Paleartic		
<i>Macrosiphoniella</i> <i>artemisiae</i> (Boyer de Fonscolombe)	NB, 1950 (ACNB)	Paleartic		
<i>Macrosiphoniella</i> <i>grandicauda</i> Takahashi and Moritsu	MD, 1997 (USNM)	Paleartic		
<i>Macrosiphoniella</i> <i>leucanthemi</i> (Ferrari)	PA, 1948 (USNM)	Paleartic		<i>Leucanthemum</i> <i>vulgare</i>
<i>Macrosiphoniella</i> <i>millefolii</i> (De Geer)	CT, 1937 (EMEC)	Paleartic		
<i>Macrosiphoniella</i> <i>sanborni</i> (Gillette)	NE, 1891 (USNM)	Paleartic (e Asia)	pest	Asteraceae (esp. chrysanthem- ums)
<i>Macrosiphoniella</i> <i>subterranea</i> (Koch)	PA, 1948 (USNM)	Paleartic		cultivated chrysanthem- ums
<i>Macrosiphoniella</i> <i>tanacetaria</i> (Kaltenbach)	CT, 1909 (CNC-Patch)	Paleartic	minor pest	cultivated chrysanthem- ums (occa- sionally)
<i>Macrosiphum lilii</i> (Monell) (19)	NY, 1878 (Comstock 1880)	Paleartic?	pest	Cultivated lilies
<i>Macrosiphum rosae</i> (Linnaeus)	MA, 1841 (Harris 1841)	Paleartic (Europe)	pest	Rosaceae (esp. <i>Rosa</i>)
<i>Macrosiphum stellariae</i> Theobald	BC, 1980 (Forbes and Chan 1983)	Paleartic	pest	Caryophyllaceae (esp. <i>Dianthus</i>)
<i>Melanaphis bambusae</i> Fullaway	GA, 1914 (USNM)	Paleartic (e Asia)		bamboos
<i>Melanaphis sacchari</i> (Zehntner)	FL, 1977 (FSCA)	Oriental	pest White et al. 2001	Gramineae (esp. sugar- cane)
<i>Melanaphis sorini</i> Halbert and Rемаудиере	FL via CA, 1996 (Halbert and Remaudière 2000)	Oriental Halbert and Remaudière 2000	pest Halbert and Remaudière 2000	<i>Miscanthus</i>
<i>Metopolophium</i> <i>dirhodum</i> (Walker)	UT, 1910 (USNM)	Paleartic (Europe)	major pest	Gramineae, incl. cereals; <i>Rosa</i>
<i>Metopolophium festucae</i> (Theobald)	CA, 1970 (USNM, Eastop 1971)	Paleartic (Europe)	pest	Gramineae
<i>Microlophium carnosum</i> (Buckton) (20)	NE, 1889 (Williams 1911)	Paleartic (Europe)		<i>Urtica</i>
<i>Microlophium sibiricum</i> (Mordvilko) (20)	PA, 1963 (USNM, Pepper 1965)	Paleartic (Europe)		<i>Urtica</i>

Table 1. Continued.

Introduced Species	Earliest N.A. record (1) (Collection or Reference)	Geographic Origin/ Reference (2)	Status in North America/ Reference (2)	North American Economic Host(s)
<i>Mindarus abietinum</i> Koch (21)	IL 1879 (Thomas 1879 [as <i>Schizoneura</i> <i>pinicola</i>])	Palaearctic (Europe)		<i>Abies</i>
<i>Monaphis antennata</i> (Kaltenbach)	ON, 1996 (CNC)	Palaearctic (Europe)		<i>Betula</i>
<i>Muscaphis musci</i> Börner	NC, 1962 (Smith and Knowlton 1975)	Palaearctic (Europe)		
<i>Myzaphis bucktoni</i> Jacob	ME, 1959 (USNM)	Palaearctic (Europe)		<i>Rosa</i>
<i>Myzaphis rosarum</i> (Kaltenbach)	NY, 1906 (USNM)	Palaearctic (Europe)	pest	<i>Rosa</i>
<i>Myzocallis carpini</i> (Koch)	BC, 1964 (Forbes et al. 1973)	Palaearctic (Europe)		<i>Carpinus</i> <i>betulus</i>
<i>Myzocallis castanicola</i> Baker	CA, 1914 (USNM)	Palaearctic (Europe)	pest	<i>Castanea</i> and <i>Quercus</i>
<i>Myzocallis coryli</i> (Goeze)	CA, 1903 (Clarke 1903)	Palaearctic (Europe)	pest	<i>Corylus</i> <i>avellana</i>
<i>Myzus antirrhini</i> (Macchiati)	MD, 1996 (USNM)	Palaearctic (Europe)		polyphagous
<i>Myzus ascalonicus</i> Doncaster	NB, 1954 (MacGillivray 1954b)	unknown Blackman and Eastop 2000	pest	polyphagous
<i>Myzus cerasi</i> Passerini	NY 1855 (Thomas 1879)	Palaearctic (Europe)	pest	<i>Prunus</i> (s.g. <i>Cerasus</i>)
<i>Myzus certus</i> (Walker)	NB, 1952 (MacGil- livray 1954a)	Palaearctic (Europe)	minor pest	Caryophyllaceae, incl. <i>Dianthus</i>
<i>Myzus hemerocallis</i> (Schrank)	FL, LA, 1958 (USNM)	Palaearctic (e Asia)	pest	<i>Hemerocallis</i>
<i>Myzus ligustri</i> (Mosley)	NY, 1915 (USNM)	Palaearctic (Europe)	occasionally severe pest Mason 1940	<i>Ligustrum</i> sp.; <i>Euonymus</i> sp.
<i>Myzus lythri</i> (Schrank)	DC, 1916 (USNM)	Palaearctic (Europe)	potential weed biocontrol agent Voegtlin 1995b	<i>Lythrum</i> <i>salicariae</i>
<i>Myzus ornatus</i> Laing	CA, 1936 (Essig 1938)	Palaearctic	pest	polyphagous
<i>Myzus persicae</i> (Sulzer)	MA, 1841 (Harris 1841)	Palaearctic (e Asia)	major pest	polyphagous
<i>Myzus varians</i> Davidson	CA 1911 (Davidson 1912)	Palaearctic (e Asia)	pest	<i>Clematis</i>
<i>Nasonovia ribisnigri</i> (Mosley)	IL, 1879 (Thomas 1879)	Palaearctic (Europe)	major pest; regulatory concern	<i>Lactuca sativa</i>
<i>Neophyllaphis araucariae</i> Takahashi	FL, 1963 (Woodruff 1963)	Australasian (Australia)	pest Russell 1967	<i>Araucaria</i> <i>heterophylla</i>
<i>Neophyllaphis podocarpi</i> Takahashi	CA, 1954 (Russell 1967)	Australasian (Australia)	pest Russell 1967, Denmark 1969	<i>Podocarpus</i>
<i>Neotoxoptera formosana</i> (Takahashi)	CA, 1927 (Essig 1935)	Oriental	pest	<i>Allium</i>
<i>Neotoxoptera oliveri</i> (Essig)	CA, 1927 (Essig 1935)	Oriental	pest	<i>Allium</i>
<i>Ovatus crataegarius</i> (Walker)	OR, 1893 (USNM)	Palaearctic (Europe)	pest	<i>Mentha</i> ; <i>Crataegus</i>
<i>Paracolopha morrisoni</i> (Baker)	MD 1916 (Baker 1919)	Palaearctic (e Asia)		bamboos
<i>Panaphis juglandis</i> (Goeze)	OR, 1928 (USNM)	Palaearctic (e Asia)	pest	<i>Juglans regia</i>

Table 1. Continued.

Introduced Species	Earliest N.A. record (1) (Collection or Reference)	Geographic Origin/ Reference (2)	Status in North America/ Reference (2)	North American Economic Host(s)
<i>Pemphigus bursarius</i> (Linnaeus)	NY, 1872 (Fitch 1872)	Palaearctic (Europe)	pest	<i>Populus nigra</i> ; <i>Lactuca</i> <i>sativa</i> <i>Populus nigra</i>
<i>Pemphigus spyrothecae</i> Passerini	QC, 1972 (Alleyne and Morrison 1974)	Palaearctic (Europe)		<i>Populus nigra</i>
<i>Pentalonia nigronervosa</i> Coquerel	DC, 1909 (USNM)	Unknown (either Neotropical or Oriental)	pest	<i>Musa</i> (banana); ornamental Musaceae, Araceae, Zingiberaceae
<i>Periphyllus californiensis</i> (Shinji)	WA, 1903 (Essig and Abernathy 1952)	Palaearctic (e Asia) Essig and Abernathy 1952		<i>Acer</i>
<i>Periphyllus lyropictus</i> (Kessler)	VT, 1847 (Fitch notes [as <i>Aphis aceris</i> L.])	Palaearctic (Europe)		<i>Acer</i> <i>platanoides</i>
<i>Periphyllus</i> <i>testudinaceus</i> (Ferne)	BC, 1924 (Essig and Abernathy 1952)	Palaearctic (Europe)		<i>Acer</i>
<i>Phloeomyzus passerini</i> (Signoret) (22)	NB? 1950, ME 1970 (Smith 1974a [as <i>P.</i> <i>dearborni</i> Smith])	Palaearctic		
<i>Phorodon humuli</i> (Schränk)	NY, 1863 (Walsh 1863)	Palaearctic (Europe)	pest	<i>Prunus</i> ; <i>Humulus</i>
<i>Phyllaphis fagi</i> (Linnaeus)	IL, 1879 (Thomas 1879)	Palaearctic (Europe)	pest	<i>Fagus</i>
<i>Pleotrichophorus</i> <i>chrysanthemi</i> (Theobald)	CA, 1910 (USNM)	Palaearctic (Asia?)	minor pest	chrysanthemums
<i>Pleotrichophorus</i> <i>glandulosus</i> (Kaltenbach)	CO, 1915 (USNM)	Palaearctic (Europe)		<i>Ambrosia</i> , <i>Artemisia</i>
<i>Prociphilus xylostei</i> (De Geer) (23)	ME, 1914 (Patch 1917)	Palaearctic		<i>Lonicera</i> ; roots of <i>Picea</i>
<i>Pseudacaudella rubida</i> (Börner)	NC, TN, 1973 (Smith and Knowlton 1975)	Palaearctic (Europe)		
<i>Pseudoregma panicola</i> (Takahashi)	FL, 1928 (FSAC)	Oriental		
<i>Pterocallis alni</i> (De Geer)	IL, 1910 (Davis 1910)	Palaearctic (Europe)		
<i>Pterocomma pilosum</i> Buckton	UT, 1958 (Knowlton 1958)	Palaearctic		
<i>Pterocomma populeum</i> (Kaltenbach)	MT, 1904 (USNM)	Palaearctic (Europe)		
<i>Pterocomma salicis</i> (Linnaeus)	MN, 1886 (Oestlund 1886)	Palaearctic		
<i>Rhopalomyzus lonicerae</i> (Siebold)	VA, 1912 (USNM)	Palaearctic (Europe)	minor pest	<i>Lonicera</i>
<i>Rhopalosiphoninus</i> <i>hydrangeae</i> (Matsumura)	BC, 1975 (Forbes and Chan 1976)	Palaearctic (e Asia)		
<i>Rhopalosiphoninus</i> <i>latysiphon</i> (Davidson)	CA, 1912 (USNM)	Palaearctic	minor pest	Bulbs; <i>Solanum</i> <i>tuberosum</i>

Table 1. Continued.

Introduced Species	Earliest N.A. record (1) (Collection or Reference)	Geographic Origin/ Reference (2)	Status in North America/ Reference (2)	North American Economic Host(s)
<i>Rhopalosiphoninus staphyleae</i> (Koch)	BC, 1953 (ACNB)	Paleartic (Europe)	minor pest	Polyphagous on roots, including stored bulbs and rootcrops
<i>Rhopalosiphum maidis</i> (Fitch) (24)	NY, 1851 (Fitch 1856)	Paleartic (e Asia)	serious pest	Gramineae
<i>Rhopalosiphum nymphaeae</i> (Linnaeus)	NE, 1890 (USNM)	Paleartic Davidson 1917	pest/ biological control	various aquatic plants
<i>Rhopalosiphum rufiabdominale</i> (Sasaki) (24)	NY, 1856 (Fitch 1856)	Paleartic (e Asia)	pest	Gramineae
<i>Schizaphis graminum</i> (Rondani)	MD, 1884 (Pergande 1902)	Paleartic	serious pest	Gramineae
<i>Schizaphis minuta</i> (van der Goot)	FL, 1998 (Halbert et al. 2001)	Oriental		
<i>Schizaphis palustris</i> (Theobald)	BC, 1974 (Forbes and Chan 1976)	Paleartic (Europe)		
<i>Schizaphis rotundiventris</i> (Signoret)	FL, 1988 (Halbert et al. 2000)	Paleartic (e Asia) or Oriental		
<i>Schizolachnus pineti</i> (Fabricius)	BC, 1961 (CNC)	Paleartic		
<i>Shivaphis celti</i> Das (25)	GA, SC, 1996 (Halbert et al. 2000)	Paleartic (E Asia) and Oriental	nuisance pest Halbert and Choate 1999	<i>Celtis</i>
<i>Sipha elegans</i> Del Guercio	NB, 1949 (ACNB)	Paleartic Black- man & Eastop 2000	minor pest	Gramineae
<i>Sipha glyceriae</i> (Kaltenbach)	ME, 1909 (Patch 1910)	Paleartic	pest	Gramineae (incl. rice)
<i>Sitobion avenae</i> (Fabricius)	NY, 1851 (Davis 1914)	Paleartic	major pest Davis 1914	Gramineae
<i>Sitobion fragariae</i> (Walker)	CA 1916 (USNM)	Paleartic (Europe)	pest	Gramineae, <i>Rubus</i>
<i>Sitobion luteum</i> (Buckton) (26)	FL, 1940 (FSCA)	unknown	minor pest	greenhouse orchids
<i>Smynthurodes betae</i> Westwood	DC, 1895 (USNM)	Paleartic (Mediterranean)	minor pest Stoetzel et al. 1996	polyphagous on roots incl. <i>Phaseolus</i> , <i>Solanum</i> , <i>Gossypium</i>
<i>Subacyrthosiphon cryptobium</i> Hille Ris Lambers	IA, 1966 (USNM)	Paleartic (Europe)		
<i>Takecallis arundicolens</i> (Clarke)	CA, 1903 (USNM)	Oriental		bamboos
<i>Takecallis arundinariae</i> (Essig)	CA, 1911 (Essig 1917)	Oriental		bamboos
<i>Takecallis taiwanus</i> (Takahashi)	FL, 1930 (USNM)	Oriental		bamboos
<i>Tetraneura fusiformis</i> Matsumura (27)	FL, 1959 (FSCA, USNM)	Paleartic (e Asia) / Oriental	possible pest	<i>Ulmus</i> ?; roots of Gramineae inc. rice
<i>Tetraneura ulmi</i> (Linnaeus)	NE, 1890 (USNM)	Paleartic		
<i>Thecabius lysimachiae</i> Börner (28)	NY, 2000 (FSCA)	Paleartic (Europe)		<i>Populus</i> , <i>Lysimachia</i>

Table 1. Continued.

Introduced Species	Earliest N.A. record (1) (Collection or Reference)	Geographic Origin/ Reference (2)	Status in North America/ Reference (2)	North American Economic Host(s)
<i>Therioaphis richmi</i> (Börner)	KS, WA, 1948 (USNM)	Palaearctic (Europe)		<i>Trifolium</i>
<i>Therioaphis trifolii</i> (Monell) (29)	NY, 1879 (USNM)	Palaearctic (Europe)	pest	Leguminosae, esp. <i>Trifolium</i> and <i>Medicago</i>
<i>Tinocallis</i> <i>kahawatuokalani</i> (Kirkaldy)	LA, 1925 (USNM)	Oriental	pest Mizell et al. 2002	<i>Lagerstroemia</i> <i>indica</i>
<i>Tinocallis platani</i> (Kaltenbach)	BC, 1927 (Glendenning 1929)	Palaearctic (Europe)		<i>Ulmus</i>
<i>Tinocallis saltans</i> (Nevsky)	ID, 1986 (Halbert and Pike 1990)	Palaearctic (e Asia)		<i>Ulmus</i>
<i>Tinocallis takachihoensis</i> Higuchi	MD, 1996 (USNM)	Palaearctic (e Asia)		<i>Ulmus</i>
<i>Tinocallis ulmiparvifoliae</i> Matsumura	CA, FL, 1998 (Gill 1998b, Halbert 1998)	Palaearctic (e Asia)	nuisance pest	<i>Ulmus</i>
<i>Tinocallis zelkowae</i> (Takahashi)	CA, 1983 (Kono 1983)	Palaearctic (e Asia)		
<i>Toxoptera aurantii</i> (Boyer de Fonscolombe)	CA, 1907 (USNM)	Palaearctic (e Asia) or Oriental	major pest	polyphagous
<i>Toxoptera citricida</i> (Kirkaldy)	FL, 1995 (Halbert and Brown 1996)	Palaearctic (e Asia) or Oriental	major pest Halbert and Brown 1996	<i>Citrus</i>
<i>Trama rara</i> Mordvilko	CO, 1929 (USNM)	Palaearctic (Europe)		roots of Compositae
<i>Trichosiphonaphis</i> <i>polygoni</i> (van der Goot)	FL, 1974 (Smith and Denmark 1982)	Palaearctic (e Asia)		<i>Polygonum</i> <i>punctatum</i>
<i>Tuberculatus annulatus</i> (Hartig) (30)	WA, 1951 (USNM)	Palaearctic (Europe)		
<i>Tuberculatus querceus</i> (Kaltenbach)	BC, 1999 (CNC)	Palaearctic (Europe)		
<i>Tuberolachmus salignus</i> (Gmelin)	IL, 1872 (Le Baron 1872)	Palaearctic (Europe)	minor pest	<i>Salix</i>
<i>Uroleucon achilleae</i> (Koch)	CA, 1966 (Hille Ris Lambers 1966)	Palaearctic (Europe)		<i>Achillea</i>
<i>Uroleucon cirsii</i> (Linnaeus)	ON, 1956 (CNC)	Palaearctic (Europe)		<i>Cirsium</i>
<i>Uroleucon sonchi</i> (Linnaeus)	DC, 1892 (USNM)	Palaearctic (Europe)		<i>Sonchus</i>
<i>Uroleucon taraxaci</i> (Kaltenbach)	NY, 1902 (USNM)	Palaearctic (Europe)		<i>Taraxacum</i>
<i>Vesiculaphis caricis</i> (Fullaway)	VA, 1916 (USNM)	Palaearctic (e Asia)		
<i>Wahlgreniella vaccinii</i> (Theobald)	IL, 1929 (USNM)	Palaearctic (Europe)		

1. Names of Canadian provinces and U.S. states indicated using standard two-letter postal abbreviations (Canada Post 2005, United States Postal Service 1998).
2. Area of origin and pest status without explicit references are based on Blackman and Eastop (1994, 2000) supplemented by opinion of the authors.
3. *Acyrtosiphon malvae*: Also known as *A. pelargonii*. This name applies to a complex of entities with various host associations. We have not distinguished records for the various forms.
4. *Aphidounguis mali*: Single collection from botanical garden.
5. *Aphis craccivora*: Previously infrequent on alfalfa, the recent upsurge of this species on alfalfa may represent a separate introduction of an alfalfa-feeding population from Asia comparable to the situation in *Therioaphis trifolii* (*A. craccivora* is known as the alfalfa aphid in China).

Table 1. Continued.

6. *Aphis euphorbiae*: Listed in Williams (1911) from NE: four slides in USNM, apparently from this collection, are not *A. euphorbiae*.
7. *Aphis fabae* and *Aphis rumicis*: The latter name was used in the past for the aphid now known as *A. fabae*. The earliest occurrence of true *A. rumicis* is thus uncertain. *Aphis fabae* is currently used for a complex of taxa usually considered as subspecies. No attempt has been made here to distinguish the members of this complex. See also under '*A. evonymi*' in Table 2.
8. *Aphis polygonata*: Single record from flight intercept trap.
9. *Aphis taraxiacicola*: Single record from trap.
10. *Brachycaudus rumexicolens*: Considered by Holman (1965) to be introduced into Europe from North America. However, the only recorded host, *Rumex acetosella*, is a Palearctic species naturalized in North America (Scoggan 1978).
11. *Dysaphis crataegi*: A number of similar *Dysaphis* species use *Crataegus* as primary host. Early reports of *D. crataegi* (e.g., Thomas 1879) based on association with *Crataegus* are not verifiable.
12. *Dysaphis foeniculus*: Single record.
13. *Eriosoma grossulariae* and *E. ulmi*: These names were long considered synonyms, but Danielsson (1982) showed that two species were involved. Reexamination of older North American collections named '*ulmi*' has shown that most, if not all, earlier collections are *E. grossulariae*. The 2003 collection of *E. ulmi* in CNC is currently the earliest confirmed record of this species in North America.
14. *Euceraphis betulae* and *E. punctipennis*: These species have been confused in the past and the correct identity of specific records is uncertain. Also, *E. betulae*, has been used to cover a complex of related non-European taxa. Most records of *E. betulae* and *E. punctipennis* from *Betula papyrifera* are actually specimens of the recently recognized species *E. papyrifericola* Blackman and subarctic records from dwarf birches probably *E. borealis* Blackman.
15. *Geocica utricularia*: Probably represents a complex of species (Blackman and Eastop 2000).
16. *Hyalopterus amygdali* and *H. pruni*: There is confusion of these two species in North America, and the exact status of individual records is uncertain.
17. *Hyperomyzus rhinanthi*: Distribution is alpine and northern Europe, Iceland, and northeastern North America. The North American population may represent part of a natural transatlantic distribution, rather than a human-mediated introduction.
18. *Lipaphis pseudobrassicae*: Records from past few decades as *L. erysimi* (Kaltenbach).
19. *Macrosiphum lili*: First described from bulbs imported from Japan. However, there have been no subsequent Japanese records (Blackman and Eastop 2000).
20. *Microlophium*: This genus is in need of revision and it is difficult to say with clarity which species are present in North America.
21. *Mindarus abietinus*: Voegtlin (1995a) believes that North American populations may represent a different species from Eurasian populations. The numerous records of this species from spruce are almost all *M. obliquus*.
22. *Phloeomyzus passerini* (Signoret): Smith (1974a) refers to a collection by N.R. Brown from "Canada" in 1950. This particular collection has not been located, but other collections by Brown in ACNB from this period are all from NB.
23. *Prociphilus xylostei*: Some early North American records of *P. xylostei* were actually based on specimens of *Hyadaphis foeniculi* (= *Aphis xylostei* Schrank) misfiled in USNM.
24. *Rhopalosiphum maidis* and *R. rufiabdominale*: Both species are assumed to be of Asian origin because complete holocyclic lifecycles are documented only in Asia. Holocyclic colonies of *R. maidis* were found on *Prunus cornuta* in Pakistan (Remaudière and Naumann-Etienne 1991). Similarly, holocyclic colonies of *R. rufiabdominale* occur on *Prunus* in Japan (Blackman and Eastop 2000). However, a holocycle for *R. rufiabdominale* also may occur in North America. (Al-Raeesi et al. 1992). There is a great deal of genetic diversity in *R. maidis* populations in North America (Blackman et al. 1990). This genetic diversity may point to a North American origin for the species, or it may be the result of mutations accumulating in continuous clonal lineages. Most other *Rhopalosiphum* species, including *Rhopalosiphum padi* (L.), are thought to be of North American origin (Halbert and Voegtlin 1998).
25. *Shivaphis celti*: Probably bonsai introductions.
26. *Sitobion luteum*: Earlier North American records in USNM represent port interceptions.
27. *Tetraneura fusiformis*: Records as *T. nigriabdominalis* (Sasaki). See Eastop and Blackman (2005).
28. *Thecabius lysimachiae*: A 1956 collection by Hille Ris Lambers in Montreal, QC was questionably assigned to this species by Hille Ris Lambers (Smith 1974b). Recent collections in New England lend credibility to his determination.
29. *Therioaphis trifolii*: Eastern populations feed on clover. Alfalfa feeding populations in western USA represent a separate introduction formerly known as *Therioaphis maculata* (or *Pterocallidium maculatum*).
30. *Tuberculatus annulatus*: Record by Lintner (1889) (as *Aphis quercus*) not verified.

Table 2. Species previously reported as introduced in North America, but now considered to be erroneous or dubious records.

Species	Reason for Consideration as Erroneous or Dubious
<i>Amphorophora rubi</i> (Kaltenbach)	May not occur in North America. References to this species on raspberry in North America are mostly misidentifications of <i>Amphorophora agathonica</i> Hottes (Blackman and Eastop 2000). However, Hille Ris Lambers believed <i>A. rubi</i> was present in northwestern North America (Hille Ris Lambers letter to L. Russell, 13 Nov. 1972, USNM letter file). Note that the European aphid on raspberry is actually <i>A. idaei</i> (Börner) while <i>A. rubi</i> is restricted to blackberry (Blackman et al. 1977).
<i>Anuraphis farfarae</i> (Koch)	Listed in Bodenheimer and Swirski (1957) as occurring in North America but no supporting specimens located.
<i>Anuraphis subterranea</i> (Walker)	Cited in Jones (1929) from Colorado. No specimens are present in USNM.
<i>Aphis frangulae</i> Kaltenbach	North American records probably refer to <i>Aphis nasturtii</i> . <i>Aphis frangulae</i> is a member of a complex of taxa including <i>Aphis gossypii</i> Glover (Smith and Parron 1978, Stroyan 1984, Blackman and Eastop 2000). The overwintering host for <i>A. frangulae sensu stricto</i> is <i>Frangula alnus</i> (= <i>Rhamnus frangula</i>). <i>A. nasturtii</i> (as <i>A. rhamni</i> Boyer de Fonscolombe, see note below for <i>Macchiatella rhamni</i>) is the only <i>Aphis</i> reported on <i>Rhamnus</i> in Illinois by Hottes and Frisson (1931).
<i>Aphis acetosae</i> Linnaeus	Listed in Börner and Heinze (1957) as occurring in North America, but no specimens have been located
<i>Aphis evonymi</i> Fabricius	There is considerable confusion in the application of the name <i>A. evonymi</i> (or <i>euonymi</i>). It is probable that most, if not all, of the North American records of <i>A. evonymi</i> (or <i>A. euonymi</i>) refer to other members of the <i>Aphis fabae</i> complex. Eastop and Hille Ris Lambers (1976) state that <i>A. evonymi</i> of authors prior to 1950 is a synonym of <i>Aphis fabae solanella</i> Theobald. Eastop and Hille Ris Lambers (1976) listed <i>A. euonymi</i> Fabricius 1775 as a valid species, and Remaudière and Remaudière (1997) treat it as a subspecies of <i>A. fabae</i> .
<i>Aphis medicaginis</i> Koch	North American records of this species are misidentifications of <i>Aphis craccivora</i> .
<i>Aphis grossulariae</i> Kaltenbach	Listed in Wilson (1913) as found in Oregon. No specimens are present in USNM.
<i>Aphis ilicis</i> Kaltenbach	Reported from California and British Columbia in Smith & Parron (1978). No supporting specimens have been located.
<i>Aphis hieracii</i> Schrank	Listed in Rafinesque (1818). No North American specimens have been located.
<i>Aphis nepetae</i> Kaltenbach	Listed in Smith and Parron (1978) as occurring in Idaho and Minnesota. There are no North American records in the CNC, FSCA, or USNM.
<i>Aphis plantaginis</i> Goeze	Listed in Smith and Parron (1978) as occurring in New York. No North American specimens have been located
<i>Aphis ulmariae</i> Schrank	Listed in Smith and Parron (1978) from various states. Specimens from Colorado in USNM were actually <i>A. spiraeicola</i> . Record by Thomas (1879) is <i>A. pisum</i> .
<i>Aphis viburni</i> Scopoli	Recorded in Illinois as early as 1879 (Thomas 1879). However, records are likely the result of confusion with <i>A. fabae</i> and <i>A. viburniphila</i> Patch.
<i>Aphis vitis</i> Scopoli	Listed in Walsh (1863) from Illinois and Missouri. A single slide of poor quality found in USNM identified as <i>A. vitis</i> is likely <i>Aphis illinoensis</i> .
<i>Brachycaudus prunicola</i> (Kaltenbach)	Listed in Hunter (1901) as occurring in North America. There are no North American records for this species in the CNC, FSCA, or USNM. Blackman and Eastop (2000) state that this species is “not yet recorded outside Europe.”

Table 2. Continued.

Species	Reason for Consideration as Erroneous or Dubious
<i>Capitophorus archangelskii</i> Nevsky	Single record from New York in 1960 (Leonard 1963). Probably <i>C. pakansus</i> . Börner and Heinze (1957) considered <i>C. pakansus</i> a synonym of <i>C. archangelskii</i> .
<i>Cinara piceicola</i> (Cholodkovsky)	North American records refer to <i>Cinara pilicornis</i> (Hartig)
<i>Cinara piceae</i> (Panzer)	No North American specimens found.
<i>Cinara pini</i> (Linnaeus)	North American records are doubtful. See Smith and Parron (1978).
<i>Cinara pinicola</i> (Kaltenbach)	No authentic North American specimens found. Records of this species appear to refer to <i>C. pinea</i> and <i>C. pilicornis</i> .
<i>Coloradoa angelicae</i> (Del Guercio)	A single published record from New York in 1913 (Olsen 1921) represents the only record of this species in North America. There are no North American specimens in CNC, FSCA, or USNM.
<i>Cryptaphis poae</i> (Hardy)	Recorded as occurring in Colorado and Manitoba (Robinson and Bradley 1965), Robinson later stated (correspondence cited in Smith and Parron 1978) that he did not believe <i>C. poae</i> occurs in North America.
<i>Dysaphis angelicae</i> (Koch)	Recorded by Wilson (1909) for California. US specimens in USNM are <i>Aphis</i> species and 2 alates and alatoid nymphs of a <i>Dysaphis</i> species that key to <i>crataegitulpipaelangelicaelpyrilradicola</i> complex. The status of this species in North America is uncertain.
<i>Dysaphis sorbi</i> (Kaltenbach)	First listed by Clarke (1903) for California. Although recorded in some subsequent literature as present in North America, specimens identified in USNM as <i>D. sorbi</i> were actually <i>D. plantaginea</i> . Stroyan (1985) did not list <i>D. sorbi</i> as present in North America.
<i>Diuraphis holci</i> (Hille Ris Lambers)	Although recorded as occurring in various localities in North America (Smith and Parron 1978), <i>D. holci</i> probably does not occur in North America (Halbert et al. 1992).
<i>Eriosoma lanigerum</i> (Hausmann)	Generally considered to be native to North America. However, as early as 1841, Harris (1841) believed otherwise and cited various sources and observations to support his view. Eastop (1973) suggests the origin may be eastern Asiatic and introduced to Europe via America.
<i>Eriosoma lanuginosa</i> (Hartig)	Probably not in North America; records refer to <i>E. pyricola</i> (Blackman and Eastop 2000).
<i>Geoica lucifuga</i> (Zehntner)	Probably not in North America; Smith (1985) believed records dubious. No specimens located.
<i>Glyphina jacutensis</i> Mordvilko	Listed as <i>G. alni</i> (Schränk) in Archibald (1957). Record is doubtful; probably actually <i>G. longiseta</i> Richards, but specimens not present in Archibald collection (now in CNC).
<i>Lachmus roboris</i> (Linnaeus)	Smith and Parron (1978) record from California, Connecticut, and Maine. No North American specimens have been located; material seen from these states identified as <i>L. roboris</i> is <i>Longistigma caryae</i> .
<i>Macchiatella rhamni</i> (Boyer de Fonscolombe)	Many North American records, but <i>Aphis rhamni</i> Boyer de Fonscolombe has been misapplied in the past by many authors to the species now called <i>Aphis nasturtii</i> Kaltenbach (Eastop and Hille Ris Lambers 1976). The presence of <i>M. rhamni</i> in North America is doubtful.
<i>Macrosiphum funestum</i> (Macchiat)	Recorded by Burnham (1938) for New Brunswick in 1936. No slides have been located in (CNC) or ACNB.
<i>Macrosiphum gei</i> (Koch)	Recorded from various localities in North America (Smith and Parron 1978). All North American specimens in FSCA and USNM were actually <i>Macrosiphum euphorbiae</i> . Blackman and Eastop (2000) state that most North American records refer to <i>M. euphorbiae</i> .
<i>Maculolachmus submacula</i> (Walker)	North American records probably all refer to <i>M. sijpkensis</i> Hille Ris Lambers (Blackman and Eastop 2000)
<i>Megoura viciae</i> Buckton	Listed in Smith and Parron (1978), citing Müller (1969). However, Müller (1969) actually states "NAs" while he uses "NAm" for North America.

Table 2. Continued.

Species	Reason for Consideration as Erroneous or Dubious
<i>Metopolophium friscium</i> Hille Ris Lambers	Listed in Smith and Parron (1978), citing Börner (1952). However, no specimens of this species have been located in North American collections.
<i>Myzus ajugae</i> Schouteden	Smith and Parron (1978) considered the Leonard (1968) reference to “ <i>Myzus ajugae</i> ” suspect, believing “the determination was made because of the host association.”
<i>Myzus myosotidis</i> (Börner)	Smith and Parron (1978) cite Hille Ris Lambers (1966) as the basis for the occurrence of <i>M. myosotidis</i> in California. However, Hille Ris Lambers (1966) mentions this species only in a key. No specimens are present in USNM. Halbert et al. (1995) considered North American distribution doubtful.
<i>Nasonovia nigra</i> (Hille Ris Lambers)	Recorded by Maw et al. (2000) in British Columbia and New Brunswick, based on mididentified specimens of <i>N. ribisnigrae</i> in ACNB.
<i>Picturaphis brasiliensis</i> (Moreira)	Cited in Smith and Parron (1978) as occurring in Florida. Examination of specimens in USNM showed the collection to be from “Florida, P.R.” (Puerto Rico), not Florida, U.S.A.
<i>Pachypappa tremulae</i> (Linnaeus)	North American records probably refer to the native counterpart, <i>P. rosettei</i> (Maxson).
<i>Pemphigus borealis</i> Tullgren	A single record from Idaho in 1952 (Gittins et al. 1976).
<i>Periphyllus aceris</i> (Linnaeus)	Although stated by Blackman and Eastop (1994) as introduced to North America, and recorded by various other authors, all North American collections in USNM are interceptions. Essig and Abernathy (1952) believed that reports of <i>P. aceris</i> were referable to other species and <i>P. aceris</i> had not been recorded in the Western Hemisphere.
<i>Prociphilus bumeliae</i> (Schrank)	According to Smith and Parron (1978), probably misidentifications of <i>P. americanus</i> .
<i>Saltusaphis scirpus</i> Theobald	A single record from Ohio, 1938 (Smith 1940). No North American specimens in USNM, CNC or FSCA.
<i>Schoutedenia rahumensis</i> Rübsaamen	A single specimen, cited in Palmer (1952) (as <i>Setaphis viridis</i> van der Goot) in Colorado. Not established.
<i>Semiaphis dauci</i> (Fabricius)	Records from District of Columbia and Maryland (Leonard and Bissell 1970, Smith and Parron 1978) may have been misidentification of <i>Dysaphis crataegi</i> .
<i>Sitobion luteum</i> (Buckton)	Cited in Smith and Parron (1978) as occurring in New Mexico and New York. All North American records in USNM are port interceptions and do not represent established populations.
<i>Thecabius affinis</i> (Kaltenbach)	<i>Thecabius affinis</i> is an Old World species, and <i>Th. populiconduplifolius</i> North American. They are morphologically and biologically indistinguishable, but <i>Th. affinis</i> has 2n = 38, and <i>Th. populiconduplifolius</i> has 2n = 28 chromosomes (Blackman and Eastop 1994). It is unknown whether <i>Th. affinis</i> occurs in North America.
<i>Thelaxes dryophila</i> (Schrank)	References to this species probably represent misidentification of <i>Thelaxes californica</i> (Davidson).
<i>Therioaphis ononidis</i> (Kaltenbach)	Records of <i>T. ononidis</i> from North America and Japan are misidentifications of <i>T. trifolii</i> (Monell) (Quednau 2003).
<i>Thripsaphis caricis</i> (Mordvilko)	North American records refer to <i>Thripsaphis producta</i> Gillette (Hille Ris Lambers 1974).
<i>Titanosiphon artemisiae</i> (Koch)	Questionably recorded for New York by Smith and Parron (1978), citing Hunter (1901) who lists “ <i>Nectarphora artemisiae?</i> Koch.” Probably a <i>Macrosiphoniella</i> species.
<i>Uroleucon compositae</i> (Theobald)	Listed in Hille Ris Lambers (1963) as present in “America.” Although introduced into the Caribbean, this species has not yet been found in North America.

Table 2. Continued.

Species	Reason for Consideration as Erroneous or Dubious
<i>Uroleucon muralis</i> (Buckton)	In Nebraska, according to in Williams (1911). However, Robinson (1985) examined specimens from the USNM and determined that they were <i>Uroleucon sonchellum</i> (Monell). He believed <i>U. muralis</i> does not occur in North America.
<i>Uroleucon solidaginis</i> (Fabricius)	All known North American specimens identified as this species are in fact various native species of <i>Uroleucon</i> . Robinson (1985) stated that he had "not seen authentic material of <i>solidaginis</i> from North America."
<i>Uroleucon tanacetii</i> (Linnaeus)	Cited in Smith and Parron (1978) as occurring in several localities. However, Robinson (1985) stated that he had "not seen authentic material of this species from North America."

Table 3. Aphid species considered to have a natural Holarctic or Beringian distribution.

<i>Aphis equiseticola</i> Ossiannilsson	The single previous North American record from Vancouver, BC suggested an introduction of this species otherwise known from Scandinavia. However, a collection (in CNC) of this species in 2004 from northern Québec indicates a possible circumboreal distribution.
<i>Aphis farinosa</i> Gmelin	Widespread in both Nearctic and Palearctic.
<i>Boernerina alni</i> Takahashi	Subspecies <i>alaskensis</i> Hille Ris Lambers and Hottes (originally described as a subspecies of <i>B. variabilis</i> Richards), widespread in Alaska and arctic Canada, is the North American segregate of this species otherwise known from east Asia.
<i>Cavariella aquatica</i> (Gillette and Bragg)	Widespread in both Nearctic and Palearctic, but infrequently collected.
<i>Cavariella konoii</i> Takahashi	According to Stroyan (1969), differences in host range among Old and New World populations indicate the possibility that this name covers a complex of taxa.
<i>Chaitophorus horii</i> Takahashi	Specimens from Northwest Territories match the nominate Asian subspecies, rather than the European subspecies (<i>beuthanii</i> Börner), and probably represent natural migration across the Bering Strait.
<i>Ghariesia polunini</i> Stroyan	Known from mountains of western North America and Himalayas.
<i>Maculolachnus sijpkensis</i> Hille Ris Lambers	Distributed across North America and in Mongolia (Blackman and Eastop 2000).
<i>Mindarus obliquus</i> (Cholodkovsky)	Present across boreal forest zone of North America, northern Europe, and northern Russia, but apparently not recorded in eastern Siberia.
<i>Muscaphis escherischi</i> (Börner)	According to Stekol'shtshikov and Shaposhnikov (1993), <i>Toxopterella drepanosiphoides</i> MacGillivray and Bradley (known from across Canada, and in northern Europe as subspecies <i>irae</i> Shaposhnikov) is the primary host form of <i>M. escherischi</i> .
<i>Myzodium modestum</i> (Hottes)	Widespread in both Nearctic and Palearctic.
<i>Thripsaphis cyperi</i> (Walker)	Although first North American records of this species in Alaska (USNM) and Ontario (CNC) date from 1961, it is distributed across boreal and subarctic Canada and is likely naturally circumboreal.
<i>Uroleucon simile</i> (Hille Ris Lambers)	Widespread in Palearctic. North American records (Alaska only) likely represent a natural expansion across Bering Strait.




Fig. 1. Adventive aphids. Number of adventive aphids in North America north of Mexico recorded at 30-year intervals. Two species not included due to confusion with related species.

(1997), except for changes noted by Eastop and Blackman (2005). The taxonomic status and biogeographic origins of some aphid species are difficult to determine conclusively due to ongoing taxonomic problems. We have summarized relevant information in footnotes to Table 1.

Parameters for determining a pest are as defined in D. R. Miller et al. (2002), Miller and Miller (2003), and Miller et al. (2005). That is, if statements are made that a species is a pest, causes harm, requires control measures, is economically important, or reflects some comparable statement or inference, then we consider that species a pest. Pest status entries in Table 1 without reference to other authors are either from Blackman and Eastop (1994, 2000) or our personal experience. The principal hosts listed are those of the greatest economic impact in North America (including crops, ornamental plants, trees and weeds). Species

causing damage to several plant families are listed as polyphagous.

A summary of species previously listed for North America, but for which we consider their occurrence doubtful or erroneous, and an explanation for their exclusion is provided in Table 2. Table 3 contains a compilation of those species, some of which were previously suspected as adventive to North America, which we consider to be most probably either Holarctic or Beringian in distribution.

RESULTS

Based on our findings there are 1,416 aphid species in North America north of Mexico, which comprises about 30% of the world fauna of approximately 4,700 species (Remaudière and Remaudière 1997). Table 1 provides information on 262 aphid species (18% of the North American and approximately 6% of the




Fig. 2. Adventive aphids. Number of adventive aphids in North America north of Mexico at 10-year intervals classified by biogeographic origin. Aphids of Neotropical, Afrotropical and unknown origin combined into single category ("Other"). Two species not included due to confusion with related species.

world aphid fauna) that are considered as adventive to North America.

There has been an overall increase in number of aphid introductions per year from 1791 until 1929 (Fig. 1, number of introductions plotted at 30-year intervals in order to smooth short term anomalies), and there has been an overall decrease to the present. There are, however, a number of irregularities in this general trend that appear when first records are plotted on a finer time scale (Fig. 2, number at 10-year intervals, and Fig. 3, cumulative number at 10-year intervals). Two species, *Aphis rumicis* L. and *Hyalopterus amygdali* (Blanchard), are excluded from Figs. 1 to 3 due to uncertain introduction dates because of confusion with related species.

The geographic origin of adventive aphids is illustrated in Fig. 2 and Fig. 3. The majority originated from the Pale-

arctic Region (227), followed by the Oriental (15), Neotropical (4), Australasian (2), and Afrotropical (1). Eight other species are generally distributed in east Asia, in both Oriental and Palearctic regions. The geographic origins of five species are unknown. As shown in other analyses (Sailer 1978, Miller et al. 2005), most insects adventive to North America originate in the Palearctic Region. We have further examined the rate of Asian introductions by subdividing the Palearctic Region into European, central Asian and east Asian components. Of the aphids originating from the Palearctic Region, 144 are found in Europe (and adjacent parts of north Africa and west Asia), 11 in central Asia, 25 in east Asia, and 48 are widespread in Eurasia. Original North American populations of the majority of the widespread Palearctic species likely originated from Europe.




Fig. 3. Adventive aphids. Cumulative number of adventive aphids in North America north of Mexico at 10-year intervals classified by biogeographic origin. Aphids of Neotropical, Afrotropical and unknown origin combined into single category ("Other"). Two species not included due to confusion with related species.

Of the 262 adventive aphids, 114 (about 44%) are considered to be pests. Of these, 20 (>17%) are recorded as either serious or major pests (i.e., causing extensive or long term damage).

DISCUSSION

The total number of adventive aphids in North America north of Mexico (262) is remarkably similar to the 255 adventive scale insects (Miller et al. 2005). Also, the decrease in annual number of introduced aphids since 1929 as shown in Fig. 1 is mirrored in the pattern seen in adventive Coccoidea (e.g., Miller et al. 2005). Sailer (1978) suggested that enforcement of the United States Plant Quarantine Act of 1912 has resulted in greater rate of interception and conse-

quent decrease in successful entry and establishment of insect species. There are also finer scale changes in the number of first detections (Fig. 2), apparently due to a combination of major geopolitical and economic events, resulting in real changes in rates of introduction, and the efforts of particular individuals in recording the aphid fauna. For example, the peak in the number of first records in the 1870's is almost entirely due to the work of Thomas (1879), and there is a major trough in the 1930's apparently related to the economic depression and associated decrease in trade activity, despite active work on aphids by Essig, Gillette, and others during this period.

As for insects in general (Sailer 1978, 1983) and for scale insects in particular

(Miller et al. 2005), most adventive aphids are indigenous to the Palearctic Region (especially Europe). This is to be expected, given the immigration history of the continent and the associated transport of crops and their weeds. However, European introductions have become relatively few (Fig. 2, and seen as a leveling off of the European component of the cumulative curve in Fig. 3). Asian species now account for a major proportion of new introductions, although the total number is similar to that in the early part of the 20th century (Fig. 2). Because the Asian fauna was poorly known at the time, several of these early oriental introductions were first described from populations in the U.S.A. (*Aphis spiraecola* Patch, *Atarsaphis agrifoliae* Ferris, *Euthoracaphis umbellariae* Essig, *Neotoxoptera olivieri* (Essig), *Paracolopha morrisoni* Baker, *Periphyllus californicus* (Shinji), *Takecallis arundicolens* (Clarke), *T. arundinariae* (Essig)) as was the African aphid *Aloephagus myersi* Essig.

The proportion of pests among the adventive species is much greater than the proportion among the native species. This to be expected, given that most of these introductions were enabled by human activity, and hence associated with economically important plants. A more detailed analysis of the economic impact of adventive and native species is in preparation.

ACKNOWLEDGMENTS

We thank D. Creel and J. Eibl (USDA, ARS, Systematic Entomology Laboratory, Beltsville, MD) for their assistance with reference verification for this paper. We also thank P. Mason (AAFC, Ottawa), E. Grissell (USDA, ARS, Systematic Entomology Laboratory, Washington, DC) and D. Nickle (USDA, ARS, Systematic Entomology Laboratory, Beltsville, MD) for their manuscript reviews.

LITERATURE CITED

- Alleyne, E. H. and F. O. Morrison. 1974. Some Canadian poplar aphid galls. *Canadian Entomologist* 109(3): 321–328.
- Al-Raeesi, A. A., S. E. Halbert, and J. B. Johnson. 1992. *Rhopalosiphum rufiabdominalis* (Sasaki) and *Aphis armoraceae* Cowen (Homoptera: Aphididae) confirmed on wheat in Idaho. *Pan-Pacific Entomologist* 68: 70–71.
- Anonymous. 2005. Soybean aphid in Minnesota. <http://www.soybeans.umn.edu/crop/insects/aphid/aphid.htm> [accessed 4 May 2005].
- Archibald, K. D. 1957. Report of the forest Aphidae (Homoptera) of Nova Scotia together with species and host plant lists. *Annals of the Entomological Society of America* 50: 38–42.
- Baker, A. C. 1919. An undescribed species of *Dryopea* injurious to *Phyllostachys*. (Aphididae-Hom.). *Proceedings of the Entomological Society of Washington* 21: 104–106.
- Baker, A. C. and W. M. Davidson. 1916. Woolly pear aphid. *Journal of Agriculture Research* 6: 351–360.
- Bartholomew, P. S. 1932. Six new species of aphids, with records of other species new to California. *Annals of the Entomological Society of America* 25(4): 713–729.
- Bernal, J., D. González, E. T. Natwick, J. G. Loya, R. León-López, and W. E. Bendixen. 1993. Natural enemies of Russian wheat aphid identified in California. *California Agriculture* 47: 24–28.
- Blackman, R. L. and V. F. Eastop. 1994. *Aphids on the World's Trees: An Identification and Information Guide*. CAB International, Wallingford, UK. 1,004 pp.
- . 2000. *Aphids on the World's Crops: An Identification and Information Guide*. Second edition. John Wiley & Sons Ltd., England. 466 pp.
- Blackman, R. L., V. F. Eastop, and M. Hills. 1977. Morphological and cytological separation of *Amphorophora* Buckton (Homoptera: Aphididae) feeding on European raspberry and blackberry (*Rubus* spp.). *Bulletin of Entomological Research* 67: 285–296.
- Blackman, R. L., S. E. Halbert, and T. W. Carroll. 1990. Association between karyotype and host plant in corn leaf aphid (Homoptera: Aphididae) in the northwestern United States. *Environmental Entomology* 19: 609–611.
- Bodenheimer, F. S. and E. Swirski. 1957. The Aphidoidea of the Middle East. Jerusalem. 378 pp.
- Börner, C. 1952. Europae centralis Aphides. Die Blattläuse Mitteleuropas. Namen, Synonyme, Wirtspflanzen, Generationszyklen. *Mitteilungen der Thüringischen Botanischen Gesellschaft* 3: 1–488.

- Börner, C. and K. Heinze. 1957. Aphidina Aphidoidea. Blattläuse, plantlice (aphids), pucerons (aphides). In Sorauer's Handbuch der Pflanzenkrankheiten 5(4): 1-402.
- Burnham, J. C. 1938. A contribution to a list of the Aphididae of the Maritime Provinces of Canada. Canadian Entomologist 70(9): 180-188.
- Canada Post, Postal guide. Section B, Chapter 3—Addressing. <http://www.canadapost.ca/tools/pg/manual/b03-e.asp> [accessed 20 August 2005].
- Chan, C. K., A. R. Forbes, and D. A. Raworth. 1991. Aphid-transmitted viruses and their vectors of the world. Agriculture Canada Technical Bulletin 1991-3E: 1-216.
- Clarke, W. T. 1903. A list of California Aphididae. Canadian Entomologist 35: 247-254.
- Cloutier, C., F. Bauduin, and F. W. Quednau. 1986. A survey of air-borne aphids in Québec by means of suction traps, with five new records for North America (Homoptera: Aphididae). Phytoprotection 67: 55-61.
- Comstock, J. H. 1880. Entomology. The Japan lily Aphis. The Valley Naturalist. 2(4): 49-50.
- Corpuz-Raros, L. A. and E. F. Cook. 1974. A revision of North American *Capitophorus* Van der Goot and *Pleotrichophorus* Börner (Homoptera: Aphididae). Smithsonian Contributions to Zoology 156: 143 pp.
- Danielsson, R. 1982. The species of the genus *Eriosoma* Leach having *Ribes* L., as secondary hostplant (Homoptera, Aphidoidea). Entomologica Scandinavica 13: 341-358.
- Davidson, W. M. 1912. Aphid notes from California. Journal of Economic Entomology 5: 404-413.
- . 1914. Plant-louse notes from California. Journal of Economic Entomology 7: 127-136.
- . 1917. The reddish-brown plum aphid (*Rhopalosiphum nymphaeae* Linn.). Journal of Economic Entomology 10: 350-353.
- Davis, J. J. 1909. Studies on Aphididae. I. Annals of the Entomological Society of America 1(4): 251-264.
- . 1910. A list of the Aphididae of Illinois, with notes on some of the species. Journal of Economic Entomology 3(5): 407-420.
- . 1914. The oat Aphis. Bulletin of the U. S. Department of Agriculture 112: 16 pp.
- Denmark, H. A. 1969. A podocarpus aphid, *Neophyllaphis podocarpi* Tak. Florida Department of Agriculture Division of Plant Industry Entomology Circular 84: 1 pp.
- Eastop, V. F. 1971. Keys for the identification of *Acyrtosiphon* (Hemiptera: Aphididae). Bulletin of the British Museum (Natural History) Entomology 26(1): 1-115.
- . 1973. Aphids and psyllids. In Viruses and Invertebrates. North-Holland Research Monographs. Frontiers of Biology. North Holland Publ. Co., Amsterdam-London, 31: 112-132.
- . 1975. Deciduous fruits and nuts. USDA Cooperative Economic Insect Report 25(18): 350.
- Eastop, V. F. and R. L. Blackman. 2005. Some new synonyms in Aphididae (Hemiptera: Sternorrhyncha). Zootaxa 1089: 1-36.
- Eastop, V. F. and D. Hille Ris Lambers. 1976. Survey of the World's Aphids. Dr. W. Junk, Publishers, The Hague. 573 pp.
- Essig, E. O. 1910. Aphididae of Southern California V. Pomona College Journal of Entomology 2: 335-338.
- . 1917. Aphididae of California. New species of Aphididae and notes from various parts of the state, but chiefly from the campus of the University of California, Berkeley, California. University of California Publications Technical Bulletins Entomology 1(7): 301-346.
- . 1932. A genus and species of the family Aphididae new to North America. University of California Publication on Entomology 6: 1-8.
- . 1935. California Aphididae. New cloudy-veined species. Pan-Pacific Entomologist 11(4): 156-162.
- . 1938. The ornate aphid, new to North America (*Myzus ornatus* Laing). Pan-Pacific Entomologist 14(2): 92-95.
- . 1950. A new genus and species of Aphididae on *Aloe* (Homoptera). Pan-Pacific Entomologist 26: 22-24.
- . 1952. A new genus and species of Aphidae on scotch broom in Oregon (Homoptera). Pan-Pacific Entomologist 28: 215-219.
- Essig, E. O. and F. Abernathy. 1952. The aphid genus *Periphyllus*. A systematic, biological, and ecological study. University of California Press, Berkeley and Los Angeles. 166 pp.
- Ferris, G. F. 1921. A new species in the Hormaphidinae (Hemiptera, Aphididae). Entomological News 32: 289-291.
- Fitch, A. 1856. First and second report of the noxious, beneficial and other insects, of the State of New York. Made to the State Agricultural Society, pursuant to an appropriation for this purpose from the legislature of the state 1856. 336 pp.
- . 1870. Thirteenth report of the noxious and other insects of the State of New York. Transactions of the New York State Agriculture Society 29: 495-566.

- . 1872. Fourteenth report of the noxious, beneficial and other insects of the State of New York. Transactions of the New York State Agriculture Society 30: 355–381.
- Forbes, A. R. and C. K. Chan. 1976. The aphids (Homoptera: Aphididae) of British Columbia 4. Further additions and corrections. Journal of the Entomological Society of British Columbia 73: 57–63.
- Forbes, A. R. 1980. The aphids (Homoptera: Aphididae) of British Columbia 8. Further additions. Journal of the Entomological Society of British Columbia 77: 38–42.
- . 1983. The aphids (Homoptera: Aphididae) of British Columbia 11. Further additions. Journal of the Entomological Society of British Columbia 80: 51–53.
- . 1984. The aphids (Homoptera: Aphididae) of British Columbia 12. Further additions. Journal of the Entomological Society of British Columbia 81: 72–75.
- Forbes, A. R., B. D. Frazer, and C. K. Chan. 1974. The aphids (Homoptera: Aphididae) of British Columbia 3. Additions and corrections. Journal of the Entomological Society of British Columbia 71: 43–49.
- Forbes, A. R., B. D. Frazer, and H. R. MacCarthy. 1973. The aphids (Homoptera: Aphididae) of British Columbia. 1. A basic taxonomic list. Journal of the Entomological Society of British Columbia 70: 43–57.
- Fox, L. 1977. Cole crops. Insects. Cooperative Plant Pest Report 2(10): 104.
- Gassmann, A., and L. T. Kok. 2002. Musk thistle (nodding thistle). In Van Driesche, R., S. Lyon, B. Blossey, M. Hoddle, and R. Reardon. 2002. Biological control of invasive plants in the Eastern United States. USDA Forest Service Publication FHTET-2002-04. 413 pp.
- Gill, R. J. 1998a. New state records: an aphid. California Plant Pest and Disease Report 17: 9.
- . 1998b. New state records: Chinese elm aphid. California Plant Pest and Disease Report 17: 10.
- Gillette, C. P. 1908. New species of Colorado Aphididae, with notes upon their life-habits. Canadian Entomologist 40: 17–20.
- . 1909. Plant louse notes, family Aphididae. Journal of Economic Entomology 2: 385–388.
- . 1910. Plant louse notes, family Aphididae (Continued), Plate 26. Journal of Economic Entomology 3(5): 403–407.
- . 1917a. Some Colorado species of the genus *Lachnus*. Annals of the Entomological Society of America 10: 133–144.
- . 1917b. Two new aphid genera and some new species. Canadian Entomologist 49: 193–199.
- Gittins, A. R., G. W. Bishop, G. F. Knowlton, and E. J. Parker. 1976. An annotated list of the aphids of Idaho (Homoptera: Aphididae). University of Idaho College of Agriculture Agricultural Experiment Station Research Bulletin 95: 47 pp.
- Glendenning, R. 1926. Some new aphids from British Columbia. Canadian Entomologist 58: 95–98.
- . 1929. A new Callipterine from Victoria, B. C. (Hemiptera-Aphididae). The Canadian Entomologist 61: 237–238.
- Hajek, A. E. and D. L. Dahlsten. 1986. Discriminating patterns of variation in aphid (Homoptera: Drepanosiphidae) distribution on *Betula pendula*. Environmental Entomology 15(6): 1145–1148.
- . 1987. The exotic aphids (Homoptera: Drepanosiphidae) on ornamental birch in Northern California. Pan-Pacific Entomologist 63(4): 319–323.
- Halbert, S. E. 1998. Triology. Entomology Section 37(3), www.doacs.state.fl.us/pi/enpp/98-may-jun.htm [accessed 14 July 2005].
- . 2004. The genus *Greenidea* (Rhynchota: Aphididae) in the United States. Florida Entomologist 87(2): 159–163.
- Halbert, S. E. and L. G. Brown. 1996. *Toxoptera citricida* (Kirkaldy), brown citrus aphid—Identification, biology and management strategies. Florida Department of Agriculture and Consumer Services Entomological Circular No. 374: 1–4.
- Halbert, S. E. and K. S. Pike. 1990. An Asian elm aphid (Homoptera: Aphididae) new to North America. Proceedings of the Entomological Society of Washington 92(4): 672–674.
- Halbert, S. E. and G. Remaudière. 2000. A new Oriental *Melanaphis* species recently introduced in North America (Hemiptera, Aphididae). Revue française d'Entomologie (Nouvelle Serie) 22(2–3): 109–117.
- Halbert, S. E. and P. M. Choate. 1999. An Asian woolly hackberry aphid, *Shivaphis celti* Das (Homoptera: Aphididae). Florida Department of Agriculture and Consumer Services Entomological Circular No. 392: 1–2.
- Halbert, S. E., S. J. Castle, and T. M. Mowry. 1995. Do *Myzus* (*Nectarosiphon*) spp. other than *M. persicae* pose a threat to the Idaho potato crop? American Potato Journal 72: 85–97.
- Halbert, S. E. and D. J. Voegtlin. 1998. Evidence for the North American origin of *Rhopalosiphum* and barley yellow dwarf virus, pp. 351–356. In Nieto, J. M., Nafria, J. M., and A. F. G. Dixon, eds. Aphid in natural and managed ecosystems. Universidad de León, León, Spain. 688 pp.

- Halbert, S. E., B. J. Connelly, and M.-g. Feng. 1992. *Diuraphis (Holcaphis) frequens* (Walker), on wheat, new to Idaho, and a key to North American *Diuraphis*. Pan-Pacific Entomologist 68: 52–61.
- Halbert, S. E., K. L. Hibbard, and D. J. Voegtlin. 2001. *Schizaphis minuta* (van der Goot) (Homoptera: Aphididae), new to the United States. Insecta Mundi 15: 56.
- Halbert, S. E., G. Remaudière, and S. E. Webb. 2000. Newly established and rarely collected aphids (Homoptera: Aphididae) in Florida and the southeastern United States. Florida Entomologist 83(1): 79–91.
- Haldeman, S. S. 1844. Descriptions of several species of *Aphis* inhabiting Pennsylvania. Proceedings of the Boston Society of Natural History 1: 168–169.
- Harper, A. M., J. P. Miska, G. R. Manglitz, B. J. Irwin, and E. J. Armbrust. 1978. The literature of arthropods associated with alfalfa: III. A bibliography of the Pea Aphid, *Acyrtosiphon pisum* (Harris) (Homoptera: Aphididae). University of Illinois at Urbana-Champaign, College of Agriculture, Agricultural Experimental Station, Special Publication, 89 pp.
- Harris, P. 2003. Classical biological control of weeds. Established biocontrol agent—*Aphis chloris* Koch. Foliage-sucking aphid. res2agr.ca/lethbridge/weedbio/agents/aaphchl_e.htm [accessed 13 July 2005].
- Harris, T. W. 1841. 2. Plant-lice. (Aphididae). A Report on the Insects of Massachusetts, Injurious to Vegetation. Cambridge, Massachusetts. 459 pp.
- Hayhurst, P. 1909. Observations on two species of *Hyalopterus* (Aphididae). Journal of the New York Entomological Society 17(3): 107–115.
- Hille Ris Lambers, D. 1949. Contributions to a monograph of the Aphididae of Europe. IV. The genera *Aulacorthum* Mord., 1914; *Microlophium* Mord., 1914; *Hyalopteroides* Theob., 1916; *Idiopterus* Davis, 1909; *Pentalonia* Coquerel, 1859; *Amphorophora* Buckton, 1876; *Wahlgreniella* nov. gen; *Megoura* Buckton, 1876; *Megourella* nov. gen.; *Hyperomyzus* Börner, 1933; *Nasonovia* Mordv., 1914. Temminckia 8: 182–323.
- . 1963. Notes on Surinam aphids. Landbouwproefstation in Suriname Bulletin 80: 1–5.
- . 1966. Notes on California aphids, with descriptions of new genera and new species (Homoptera: Aphididae). Hilgardia 37(15): 569–623.
- . 1974. On American aphids, with descriptions of a new genus and some new species (Homoptera, Aphididae). Tijdschrift voor Entomologie 117(4): 103–155.
- Holman, J. 1965. Some unrecorded Middle European aphids. Acta Faunistica Entomologica Musei Nationalis Pragae 11: 277–284.
- Hottes, F. C. and E. O. Essig. 1955. A new species of *Cinara* from Oregon (Aphidae). Proceedings of the Biological Society of Washington 68: 61–63.
- Hottes, F. C. and T. H. Frison. 1931. The plant lice of Illinois. Illinois Department of Registration and Education Division of the Natural History Survey Bulletin 19: 121–447.
- Huber, D. M., M. E. Hugh-Jones, M. K. Rust, S. R. Sheffield, D. Simberloff, and C. R. Taylor. 2002. Invasive pest species: Impacts on agricultural production, natural resources, and the environment. CAST Issue Paper 20: 1–18.
- Hunter, W. D. 1901. The Aphididae of North America. Iowa Agricultural College Experiment Station, Ames, Iowa. Bulletin 60: 63–138.
- Jones, C. M. 1929. Ants and their relation to aphids. Colorado Experiment Station Bulletin 341: 1–96.
- Knowlton, G. F. 1958. Some Utah insects—1958. Utah State University Extension Service Mimeograph Series 171: 1–16.
- Knowlton, G. F. and C. F. Smith. 1936. Aphid studies. Festschrift zum 60. Geburtstage von Professor Dr. Embrik Strand 1: 487–492.
- Kono, T. 1977. Distribution and identification of Blue Alfalfa Aphid *Acyrtosiphon kondoi* Shinji (Homoptera: Aphididae). Cooperative Plant Pest Report 2(11): 153–156.
- . 1983. A Japanese Zelkova aphid new to the U.S.A. California Plant Pest and Disease Report 2: 107–109.
- Le Baron, W. 1872. Insects injurious to the willow. Report of the State Entomologist (Illinois) 2: 138–139.
- Leonard, M. D. 1963. A list of aphids of New York. Proceedings of the Rochester Academy of Science 10(6): 289–432.
- . 1968. A supplement to a list of the aphids of New York. Proceedings of the Rochester Academy of Science 11: 257–361.
- . 1970. New state records. USDA Cooperative Economic Insect Report 20: 773.
- Leonard, M. D. and T. L. Bissell. 1970. A list of the aphids of District of Columbia, Maryland and Virginia. University of Maryland Agricultural Experiment Station. MP 770. College Park, Maryland. 129 pp.
- Lintner, J. A. 1889. The Aphididae, or plant-lice. Transactions New York State Agricultural Society 1883–1886: 101–115.

- MacGillivray, M. E. 1954a. Note on *Myzus certus* (Walker), an aphid new to North America (Homoptera: Aphidae). Canadian Entomologist 86(4): 190.
- . 1954b. Note on *Myzus ascalonicus* Doncaster (Homoptera: Aphidae), an aphid new to North America. Canadian Entomologist 86(10): 454.
- Mason, P. W. 1940. A revision of the North American aphids of the genus *Myzus*. U.S. Department of Agriculture Miscellaneous Publication 371, 31 pp.
- Maw, H. E. L., R. G. Foottit, K. G. A. Hamilton, and G. G. E. Scudder. 2000. Checklist of the Hemiptera of Canada and Alaska. NRC Research Press, Ottawa, Ontario, Canada. 220 pp.
- Miller, D. R., G. L. Miller, and G. W. Watson. 2002. Invasive species of mealybugs (Hemiptera: Pseudococcidae) and their threat to U.S. agriculture. Proceedings of the Entomological Society of Washington 104(4): 825–836.
- Miller, D. R., G. L. Miller, G. S. Hodges, and J. A. Davidson. 2005. Introduced scale insects (Hemiptera: Coccoidea) of the United States and their impact on U.S. Agriculture. Proceedings of the Entomological Society of Washington 107(1): 123–158.
- Miller, G. L. and D. R. Miller. 2003. Invasive soft scales (Hemiptera: Coccidae) and their threat to U. S. agriculture. Proceedings of the Entomological Society of Washington 105(4): 832–846.
- Miller, G. L., M. B. Stoetzel, R. López, and D. A. Potter. 2002. *Geocia setulosa* (Passerini) (Hemiptera: Aphididae): New distribution records for North America. Proceedings of the Entomological Society of Washington 104(1): 160–163.
- Mizell, R. F., F. D. Bennett, and D. K. Reed. 2002. Unsuccessful search for parasites of the crape-myrtle aphid, *Tinocallis kahawailokalani* (Homoptera: Aphididae). Florida Entomologist 85(3): 521–523.
- Morrison, W. P. and F. B. Peairs. 1998. Response model concept and economic impact. In Quisenberry, S. S. and F. B. Peairs, eds. Response model for an Introduced Pest—The Russian wheat aphid. Proceedings Thomas Say Publications in Entomology, Entomological Society of America, Lanham, Maryland, 442 pp.
- Müller, F. P. 1969. Homoptera Aphidina. Unterordnung Aphidina—Blattläuse, Aphiden. Exkursionsfauna von Deutschland. Wirbellose II/2: 51–141.
- Oestlund, O. W. 1886. List of the Aphididae of Minnesota, with descriptions of some new species. Annual Report Geological and Natural History Survey of Minnesota 14: 17–56.
- . 1887. Synopsis of the Aphididae of Minnesota. Geological and Natural History Survey of Minnesota Bulletin 4: 1–100.
- Olsen, C. E. 1921. Two season's collecting of Aphidae, principally on Long Island, New York, with notes on some of the species. Bulletin of the Brooklyn Entomological Society 16: 14–19.
- Olsen, W. H. 1974. Dusky-veined walnut aphid studies. California Agriculture 28: 18–19.
- Paddock, F. B. 1915. The turnip louse. Texas Agricultural Experiment Station Bulletin 180: 77 pp.
- Palmer, M. A. 1952. Aphids of the Rocky Mountain region. The Thomas Say Foundation 5: 1–452.
- Patch, E. M. 1910. Four rare aphid genera from Maine. Maine Agriculture Experiment Station Bulletin 182: 241–248.
- . 1911. Aphididae. Insect Notes for 1910. Maine Agriculture Experiment Station Bulletin 187: 20–23.
- . 1912. Food plant catalogue of the Aphidae of the world. Part I. (A bibliographical appendix to aphid pests of Maine.) Maine Agriculture Experiment Station Bulletin 202: 179–214.
- . 1914. Currant and gooseberry aphids in Maine. Maine Agriculture Experiment Station Bulletin 225: 49–60.
- . 1917. Eastern aphids, new or little known. Part I. Journal of Economic Entomology 10(4): 416–420.
- . 1918. Food plant catalogue of the Aphidae of the world. Part V. Maine Agriculture Experiment Station Bulletin 270: 47–100.
- Pergande, T. 1902. The Southern grain louse (*Toxoptera graminum* Rondani). Some miscellaneous results of the work of the division of entomology. VI. U.S. Department of Agriculture, Division of Entomology Bulletin 38: 7–19.
- Pepper, J. O. 1965. A list of the Pennsylvania Aphididae and their host plants (Homoptera). Transactions of the American Entomological Society 91: 181–231.
- Quednau, F. W. 1966. A list of aphids from Quebec with descriptions of two new species (Homoptera: Aphidoidea). Canadian Entomologist 98(4): 415–430.
- . 2003. Atlas of the Drepanosiphine aphids of the world. Part II: Panaphidini Oestlund, 1923—Panaphidina Oestlund, 1923 (Hemiptera: Aphididae: Calaphidinae). Memoirs of the American Entomological Institute 72: 1–301.
- Quisenberry, S. S. and F. B. Peairs. 1998. Response model for an introduced pest—the Russian Wheat Aphid. Proceedings Thomas

- Say Publications in Entomology. Entomological Society of America, Lanham, Maryland. 442 pp.
- Race, S. R. 1970. New state records. USDA Cooperative Economic Insect Report 20: 156.
- Rafinesque, C. S. 1818. Second memoir on the genus *Aphis*, containing the descriptions of 24 new American species. American Monthly Magazine 3: 15–18.
- Ragsdale, D. W., D. J. Voegtlin, and R. J. O'Neil. 2004. Soybean aphid biology in North America. Annals of the Entomological Society of America 97(2): 204–208.
- Remaudière, G. and K. Naumann-Etienne. 1991. Découverte au Pakistan de l'hôte primaire de *Rhopalosiphum maidis* (Fitch) (Hom. Aphididae). Comptes rendus de l'Académie d'Agriculture de France 77: 61–62.
- Remaudière, G. and M. Remaudière. 1997. Catalogue des Aphididae du monde (Homoptera Aphidoidea). INRA Editions, Paris. 473 pp.
- Riley, C. V. and J. Monell. 1879. Notes on the Aphididae of the United States, with descriptions of species occurring west of the Mississippi. Part II. Notes on Aphidinae, with descriptions of new species. Bulletin of the United States Geological and Geographical Survey of the Territories 5(1): 1–32.
- Robinson, A. G. 1985. Annotated list of *Uroleucon* (*Uroleucon*, *Uromelan*, *Satula*) (Homoptera: Aphididae) of America north of Mexico, with keys and descriptions of new species. Canadian Entomologist 117: 1029–1054.
- Robinson, A. G. and G. A. Bradley. 1965. A preliminary list of the aphids of Manitoba. Proceedings of the Entomological Society of Manitoba 21: 39–45.
- Russell, L. M. 1967. Notes on, and a key to, the species of *Neophyllaphis* Takahashi (Homoptera: Aphididae). USDA Cooperative Economic Insect Report 17(11): 187–194.
- . 1968. *Aphis gossypii* Glover 1877 (Insecta, Hemiptera-Homoptera): Request for validation under the plenary powers. Z. N. (S.). Bulletin of Zoological Nomenclature 25: 116–119.
- Sailer, R. I. 1978. Our immigrant insect fauna. Bulletin of the Entomological Society of America 24: 3–11.
- . 1983. History of insect introductions, pp. 15–38. In Wilson, C. L. and C. L. Graham, eds. Exotic Plant Pests and North American Agriculture. Academic Press, New York, 522 pp.
- Scoggan, H. J. 1978. The Flora of Canada. Part 3 Dicotylendoneae (Saururaceae to Violaceae). National Museum of Natural Sciences Publications in Botany. No. 7(3). National Museums of Canada, Ottawa. 1,115 pp.
- Smith, C. F. 1940. Notes on some Ohio aphids. Ohio Journal of Science 40(3): 139–142.
- . 1974a. The genus *Phloeomyzus* with the description of *P. dearborni* n. sp. from *Populus tremuloides* Michx. (Homoptera: Aphididae). Proceedings of the Entomological Society of Washington 76: 66–72.
- . 1974b. Keys to and descriptions of the genera of Pemphigini in North America (Homoptera: Aphididae: Pemphiginae). North Carolina Agricultural Experiment Station Technical Bulletin No. 226, 61 pp.
- . 1985. *Pemphiginae* in North America. In Szelegiewicz, H. ed. Evolution and biosystematics of aphids. Proceedings of the International Aphidological Symposium at Jablonna, 5–11 April, 1981. Polska Akademia Nauk Instytut Zoologii, 510 pp.
- Smith, C. F. and H. A. Denmark. 1982. *Trichosiphonaphis polygoni* (van der Goot) (Homoptera: Aphididae), a genus and species new to the United States. Florida Entomologist 65(3): 381–382.
- Smith, C. F. and G. F. Knowlton. 1975. Moss aphids in the United States (Homoptera: Aphididae). USDA Cooperative Economic Insect Report 25(21): 423–431.
- Smith, C. F. and C. S. Parron. 1978. An annotated list of Aphididae (Homoptera) of North America. North Carolina Agricultural Experiment Station Technical Bulletin 255: 1–428.
- Smith, E. F. 1890. The black peach aphid. A new species of the genus *Aphis*. Entomologica Americana 6: 101–103.
- Stekol'shtshikov, A. V. and G. K. Shaposhnikov. 1993. Revision of aphids of the genus *Musaphis* Börner (Homoptera: Aphididae). Entomologicheskoye Obozreniye 72(2): 333–344. (In Russian; English translation, 1994 Entomological Review 73(1): 42–54).
- Stoetzel, M. B. 1985. *Eucarazzia elegans* (Ferrari), an aphid new to the western hemisphere, with archival data (Homoptera: Aphididae). Proceedings of the Entomological Society of Washington. 87(1): 44–48.
- . 1987. Information on and identification of *Diuraphis noxia* (Homoptera: Aphididae) and other aphid species colonizing leaves of wheat and barley in the United States. Journal of Economic Entomology 80: 696–704.
- . 1990. Some aphids of importance to the southeastern United States (Homoptera: Aphididae). Florida Entomologist 73(4): 580–586.

- Stoetzel, M. B., G. L. Miller, P. J. O'Brien, and J. B. Graves. 1996. Aphids (Homoptera: Aphididae) colonizing cotton in the United States. *Florida Entomologist* 79(2): 193–205.
- Stroyan, H. L. G. 1969. Notes on some species of *Cavariella* Del Guercio, 1911 (Homoptera: Aphidoidea). *Proceedings of the Royal Entomological Society, London (B)* 38: 7–19, Plates I–II.
- . 1984. Aphids—Pterocommatinae and Aphidinae (Aphidini) Homoptera, Aphididae. *Handbooks for the Identification of British Insects*, Vol. 2, Part 6. Royal Entomological Society of London. London, 232 pp.
- . 1985. Recent developments in the taxonomic study of the genus *Dysaphis* Börner. In Szelegiewicz, H., ed. *Evolution and biosystematics of aphids*. *Proceedings of the International Aphidological Symposium at Jablonna*, 5–11 April, 1981. Polska Akademia Nauk Instytut Zoologii, 510 pp.
- Thomas, C. 1879. Noxious and beneficial insects of the state of Illinois. *Report of the State Entomologist (Illinois)* 8: 1–212.
- Tissot, A. N. 1944. Additions to the Lachnini of Florida (Homoptera: Aphididae). *Florida Entomologist* 27(3): 43–54.
- United States Postal Service, Official USPS Abbreviations. http://www.usps.com/ncsc/lookups/usps_abbreviations.html [accessed 20 August 2005].
- Voegtlin, D. J. 1981. Notes on a European aphid (Homoptera: Aphididae) new to North America. *Proceedings of the Entomological Society of Washington* 83(2): 361–362.
- . 1995a. Notes on the *Mindarus* spp. (Homoptera: Aphididae) of North America with descriptions of two new species. *Proceedings of the Entomological Society of Washington* 97(1): 178–196.
- . 1995b. Potential of *Myzus lythri* (Homoptera: Aphididae) to influence growth and development of *Lythrum salicaria* (Myrtiflorae: Lythraceae). *Environmental Entomology* 24(3): 724–729.
- . 2000. Soybean aphid discovered in Illinois. www.inhs.uiuc.edu/cbd/aphid/index.html [accessed 13 July 2005].
- Walsh, B. D. 1863. On the genera of Aphidae found in the United States. *Proceedings of the Entomological Society of Philadelphia* 1: 294–311.
- Wheeler, A. G., Jr and E. R. Hoebeke. 2001. A history of adventive insects in North America: Their pathways of entry and programs for their detection, pp. 3–15. In USDA-APHIS. *Detecting and monitoring invasive species*. *Proceedings of the Plant Health Conference October 24–25, 2000*. United States Department of Agriculture, Animal and Plant Health Inspection Service, Technical Bulletin, 79 pp.
- White, W. H., T. E. Reagan, and D. G. Hall. 2001. *Melanaphis sacchari* (Homoptera: Aphididae), a sugarcane pest new to Louisiana. *Florida Entomologist* 84(3): 435–436.
- Williams, T. A. 1911 (1910). The Aphididae of Nebraska. *University Studies (Nebraska)* 10: 85–175.
- Wilson, H. F. 1909. Some new records of Aphididae in North America. *Journal of Economic Entomology* 2: 346–350.
- . 1913. Plant lice attacking orchard and bush fruits in Oregon. *Biennial Crop Pest and Horticulture Report Oregon Agriculture College Experiment Station 1911–1912*: 81–96.
- . 1919. Some new Lachnids of the genus *Lachniella*. *Canadian Entomologist* 51(1): 18–22.
- Woodruff, R. E. 1963. New or rarely found insects. *Florida Cooperative Insect Pest Survey* 10(44): 131.